

From the Department of Medicine  
Karolinska Institutet, Stockholm, Sweden

**IDENTIFICATION AND CHARACTERIZATION OF  
NOVEL GENETIC AND EPIGENETIC FACTORS  
REQUIRED FOR NORMAL AND MALIGNANT  
HEMATOPOIESIS**

Yaser Heshmati



**Karolinska  
Institutet**

Stockholm 2018

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Eprint AB 2018

© Yaser Heshmati, 2018

ISBN 978-91-7831-164-4

# Identification and characterization of novel genetic and epigenetic factors required for normal and malignant hematopoiesis

## THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

**Yaser Heshmati**

*Principal Supervisor:*

Assistant Professor Julian Walfridsson, Ph.D.  
Karolinska Institutet  
Department of Medicine  
Center for Hematology and  
Regenerative Medicine (HERM)

*Co-supervisors:*

Assistant Professor Hong Qian, Ph.D.  
Karolinska Institutet  
Department of Medicine  
Center for Hematology and  
Regenerative Medicine (HERM)

Professor Petter Höglund, M.D., Ph.D.  
Karolinska Institutet  
Department of Medicine  
Center for Hematology and  
Regenerative Medicine (HERM)

*Opponent:*

Professor Tim Somervaille, Ph.D.  
University of Manchester  
Department of Biology, Medicine and Health  
Division of Cancer Research UK (CRUK)

*Examination Board:*

Professor Anthony Wright, Ph.D.  
Karolinska Institutet  
Department of Laboratory Medicine (LABMED)  
Division of Clinical Research Center

Professor Helena Jernberg Wiklund, Ph.D.  
Uppsala University  
Department of Immunology, Genetic and  
Pathology (IGP)  
Division of Experimental and Clinical Oncology

Assistant Professor Larry Mansouri, Ph.D.  
Karolinska Institutet  
Department of Medicine  
Division of Molecular Medicine and Surgery



The desire for science I could not forego,  
Few secrets remained that I did not know  
Seventy-two years, day and night I thought  
Yet I came to know, I have nothing to show  
**Omar Khayyam (1048-1131)**

To Scientists,  
who have always devoted their lives to a better world



## ABSTRACT

Acute myeloid leukemia (AML) is a type of blood cancer, characterized by clonal expansion and loss of differentiation ability of myeloid progenitor cells leading to abnormal accumulation of immature myeloid cells (myeloblasts) in the bone marrow and peripheral blood.

This thesis (study I to IV) focused on the identification and characterization of genes which are required for AML growth. The final study (study V) aimed to uncover the role of *NAPIL3* in normal hematopoietic stem cells (HSCs).

In studies I and II, we performed large-scale RNA interference screens in mouse and AML human cell lines to identify novel factors and pathways required for AML growth. Using this approach, we identified two novel targets: Chromatin remodeling factor *CHD4* (study I) and the transcription factor *GTF2IRD1* (study II), which display both a strong inhibitory effect on the growth of AML cells and a less negative effect on normal hematopoietic cells.

Using RNA interference and CRISPR-Cas9 techniques, we revealed that these genes were crucial for AML cell growth *in vitro* and *in vivo*. Knockdown of either *CHD4* or *GTF2IRD1* accumulated cells in the G0 phase of the cell cycle and resulted in downregulation of *MYC* and its target genes. We demonstrated the inhibitory role of *CHD4* knockdown on the growth and maintenance of primary childhood AML in an *ex vivo* setting, as well as in a xenograft model by transplanting patient-derived samples into humanized NSG-SGM3 mice. *GTF2IRD1* knockdown reduced the number of primary childhood and adult AML cells in *ex vivo* culture and delayed AML progression in the transplanted animal model. Therefore, *CHD4* and *GTF2IRD1* are important for AML cell growth, and interestingly the knockdown of these two genes did not show a strong inhibitory effect on normal hematopoietic cell growth.

In study III, we described the role of an epigenetic enzyme, the histone methyl-transferase *EHMT1* in AML. We used RNA interference, CRISPR-Cas9, and pharmacological approaches to inhibit *EHMT1* expression, which prevented the growth of various AML cell lines and primary AML patient samples. Knockdown of *EHMT1* significantly delayed disease progression in AML mouse models and prolonged their survival. Next, we employed CRISPR-Cas9 technology to generate single and double gene knockouts of *EHMT1* and its homolog *EHMT2*, which showed that both enzymes cooperatively play a role in AML cell proliferation and shared a similar cellular mechanism as individual knockouts of either gene resulted in an increased number of cells in G0 phase of the cell cycle. RNA sequencing of

the transcriptome of AML cells with *EHMT1* and *EHMT2* knockdown identified several common biological processes, including cell differentiation, proliferation and survival, as well as other unshared pathways and downstream effectors.

In study IV, we contributed to Nikolas Herold's study, who found that deoxynucleoside triphosphate (dNTP) triphosphohydrolase SAM domain and HD domain 1 (*SAMHD1*) plays a role in detoxifying intracellular ara-CTP in cells treated with the deoxycytidine analog cytarabine (ara-C). Transient reduction of *SAMHD1* expression by using the simian immunodeficiency virus (SIV) protein Vpx significantly increased the sensitivity of AML cells to ara-C, whereas AML cells lacking *SAMHD1* transplanted into recipient mice were hypersensitive to ara-C. We showed that *in vitro* treatment of primary AML patient samples with Vpx, which suppresses *SAMHD1*, resulted in reduced proliferation of AML but not normal cells. Together, our data suggest that *SAMHD1* inhibition can be used as a therapeutic strategy for cancer (AML) patients with high *SAMHD1* expression.

In study V, our aim was to identify novel epigenetic regulators of normal HSCs. We found high expression level of *Nap1l3*, a member of nucleosome assembly proteins (NAPs), as a histone chaperone in HSCs. Loss of function of mouse *Nap1l3* mediated by shRNA or CRISPR-Cas9 impaired the maintenance and differentiation of HSCs in both our *in vitro* and *in vivo* studies. Moreover, downregulation of *NAP1L3* in human UCB HSCs significantly decreased both the number of colonies formed by HSCs and their proliferation *in vitro* due to cell cycle arrest in the G0 phase. Xenograft mouse models using human HSCs with *NAP1L3* knockdown showed a reduction of HSC reconstitution and bias in differentiation. Furthermore, we observed upregulation of several *HOX* genes (*HOXA3*, *HOXA5*, *HOXA6* and *HOXA9*) under *NAP1L3* suppression in human HSCs.

Altogether, in this thesis, we showed the important roles of *CHD4*, *EHMT1* and *GTF2IRD1* in AML cell expansion, identifying them as potential novel targets for AML treatment. Moreover, we revealed the cellular mechanisms and RNA expression patterns under knockdown of these genes. We contributed to the study that found *SAMHD1* expression level can be used as a prognostic marker for ara-C treatment and that inhibition of *SAMHD1* increases the sensitivity of AML cells to ara-C treatment. Finally, we identified a novel regulatory role for *NAP1L3* as a histone chaperone in self-renewal and differentiation of HSCs.





## LIST OF SCIENTIFIC PAPERS

- I. **The chromatin-remodeling factor CHD4 is required for maintenance of childhood acute myeloid leukemia.**  
Yaser Heshmati, Gözde Türköz, Aditya Harisankar, Shabnam Kharazi, Johan Boström, Esmat Kamali Dolatabadi, Aleksandra Krstic, David Chang, Robert Månsson, Mikael Altun, Hong Qian and Julian Walfridsson. *Haematologica*, 2018, volume 103 (7), 1169-1181.
- II. **Identification of GTF2IRD1 as a novel transcription factor essential for acute myeloid leukemia.**  
Yaser Heshmati, Gözde Türköz, Marios Dimitriou, Aditya Harisankar, Johan Boström, Mikael Altun, Hong Qian, Nadir Kadri, Julian Walfridsson. *Manuscript*.
- III. **The histone methyltransferase EHMT1 plays both independent and cooperative regulatory role in the maintenance of Acute Myeloid Leukemia**  
Yaser Heshmati\*, Gözde Türköz\*, Emma Wagner, Aditya Harisankar, Johan Boström, Mikael Altun, Hong Qian and Julian Walfridsson. *Manuscript*.  
\*Authors contributed equally to this study
- IV. **Targeting SAMHD1 with the Vpx protein to improve cytarabine therapy for hematological malignancies.**  
Nikolas Herold, Sean G Rudd, Linda Ljungblad, Kumar Sanjiv, Ida Hed Myrberg, Cynthia B J Paulin, Yaser Heshmati, Anna Hagenkort, Juliane Kutzner, Brent D G Page, José M Calderón-Montaña, Olga Loseva, Ann-Sofie Jemth, Lorenzo Bulli, Hanna Axelsson, Bianca Tesi, Nicholas C K Valerie, Andreas Höglund, Julia Bladh, Elisée Wiita, Mikael Sundin, Michael Uhlin, Georgios Rassidakis, Mats Heyman, Katja Pokrovskaja Tamm, Ulrika Warpmann-Berglund, Julian Walfridsson, Sören Lehmann, Dan Grandér, Thomas Lundbäck, Per Kogner<sup>1</sup>, Jan-Inge Henter, Thomas Helleday & Torsten Schaller. *Nature Medicine*, 2017, Volume 23, 252-263.
- V. **The histone chaperone NAP1L3 is required for hematopoietic stem cell maintenance and differentiation.**  
Yaser Heshmati, Shabnam Kharazi, Gözde Türköz, David Chang, Esmat Kamali Dolatabadi, Johan Boström, Aleksandra Krstic, Theodora Boukoura, Emma Wagner, Nadir Kadri, Robert Månsson, Mikael Altun, Hong Qian, Julian Walfridsson. *Scientific reports*, 2018, volume 8(1)11202.

## **RELATED PUBLICATIONS NOT INCLUDED IN THE THESIS**

### **Distinct roles of mesenchymal stem and progenitor cells during the development of Acute Myeloid Leukemia in mice.**

Pingnan Xiao, Lakshmi Sandhow, **Yaser Heshmati**, Makoto Kondo, Thibault Boudierlique, Monika Dolinska, Anne-Sofie Johansson, Mikael Sigvardsson, Marja Ekblom, Julian Walfridsson, and Hong Qian. *Blood advances*, 2018, *Volume 2*, 1480-1494.

### **Xeno-immunosuppressive properties of human decidual stromal cells in mouse models of alloreactivity in vitro and in vivo.**

Behnam Sadeghi, **Yaser Heshmati**, Bita Khoein, Helena Kaipe, Mehmet Uzunel, Julian Walfridsson, Olle Ringden. *Cytotherapy*, 2015, *Volume 17*, 1732-1745.

# CONTENTS

1	Introduction .....	1
1.1	Hematopoietic system .....	1
1.1.1	Hematopoiesis .....	2
1.1.2	Hematopoietic stem cells and differentiation .....	3
1.1.3	Epigenetic regulation of hematopoiesis.....	9
1.1.4	Transcription factors involved in hematopoiesis.....	14
1.2	Acute Myeloid Leukemia .....	18
1.2.1	Diagnosis of AML .....	18
1.2.2	Classification of AML .....	20
1.2.3	Mutational landscape of AML.....	22
1.2.4	Prognosis factors .....	28
1.2.5	The bone marrow niche of AML.....	30
1.2.6	Clonal hematopoiesis and AML evolution.....	31
1.2.7	Leukemic stem cells .....	32
1.2.8	AML treatment .....	33
1.2.9	Childhood AML .....	39
2	Aim of the thesis .....	41
3	Methodological approaches .....	43
3.1	Lentiviral transfection and transduction.....	43
3.2	Large scale shRNA screen .....	43
3.3	CRISPR/Cas 9 genome editing .....	45
3.4	Flow cytometric analysis and sorting.....	47
3.5	Cell growth assays of AML patient samples.....	49
3.6	Isolation and culture of primary normal cells .....	49
3.7	Colony forming unit assay .....	50
3.8	AML mouse models and transplantation studies .....	50
3.9	RNA Sequencing.....	51
4	Results and discussion .....	53
4.1	Results .....	53
4.1.1	Study I .....	53
4.1.2	Study II .....	55
4.1.3	Study III.....	57
4.1.4	Study IV.....	59
4.1.5	Study V .....	60
4.2	Discussion .....	62
5	Concluding remarks .....	67
6	Acknowledgements.....	69
7	References .....	75

## LIST OF ABBREVIATIONS

5-hmC	5-hydroxymethylcytosine
5mC	5-methylcytosine
ADP	Adenosine diphosphate
AGM	Aorta-Gonad-Mesonephros
AML	Acute Myeloid Leukemia
ASXL1	Additional Sex Combs Like 1
BAD	Bcl-XL/Bcl-2-Associated Death
bHLH	basic-Helix-Loop-Helix
BM	Bone Marrow
BMI1	B Lymphoma Mo-MLV Insertion region 1
BMP4	Bone Morphogenetic Protein 4
BrdUrd	BromodeoxyUridine
C/EBP $\alpha$	CCAAT/Enhancer Binding Protein alpha
CAR	CXC chemokine ligand (CXCL)12-Abundant Reticular
CAR	Chimeric Antigen Receptor
CARTs	CAR-transduced T cells
CBFA2	Core-Binding Factor subunit $\alpha$ -2
CBL	Casitas B-Lineage Lymphoma
CCL3	C-C Motif Chemokine Ligand 3
CD	Cluster of Differentiation
CEBPA	CCAAT Enhancer Binding Protein A
CFU	Colony Forming Unit
CFU-E/BFU-E	CFU-Erythrocytes
CFU-G/GM	CFU-Granulocytes/Macrophages
CFU-GEM	CFU-Granulocyte, Erythrocyte, Monocyte/macrophage
CFU-M	CFU-Macrophages

CH	Clonal Hematopoiesis
CHD4	Chromodomain Helicase DNA Binding Protein 4
CLP	Common Lymphoid Progenitors
CMP	Common Myeloid Progenitor
CN-AML	Cytogenetically Normal AML
CR	Complete Remission
CRISPR	Clustered Regularly Interspaced Short Palindrome Repeats
CRM1	Chromosome Region Maintenance 1
CXCL12	C-X-C Motif Chemokine Ligand 12
DNMT	DNA Methyltransferase
DNMT3A	DNA Methyltransferase 3A
DOT1L	Disruptor of Telomeric silencing 1-Like
DSBs	Double Strand Breaks
dsRNA	double strand RNA
EHMT1/2	Euchromatic Histone Lysine Methyltransferase 1/2
ELN	European LeukemiaNet
EMP	Erythroid and Myeloid Progenitors
ERK1/2	Extracellular signal-Related Kinase 1/2
ETV6	ETS Variant 6
FAB	French-America-British
FACS	Fluorescent Activated Cell Sorter
FDCP	Factor-Dependent Continuous Paterson Laboratories
FcR	Fragment crystallizable Receptor
FDA	Food and Drug Administration
FISH	Fluorescent In Situ Hybridization
FLT3	Fms-Like Tyrosine kinase 3
GAB1	GRB2-Associated Binder 1

GADD45a	Growth Arrest and DNA Damage Inducible Alpha
GFI1	Growth Factor Independent 1
GFP	Green Fluorescent Protein
GMP	Granulocyte-Monocyte Progenitor
GNAT	Glycine N-Acyltransferase-Like Protein 1
GRB2	Growth Factor Receptor-Bound2
GSEA	Gene Set Enrichment Analysis
GTF2IRD1	GTF2I Repeat Domain Containing 1
GVHD	Graft Versus Host Disease
HAT	Histone AcetylTransferase
HDAC	Histone Deacetylase
HDR	Homology-Directed Repair
HES1	Hairy and Enhancer of Split 1
HLA	Human Leukocyte Antigen
HLA-DR	Human Leukocyte Antigen D-related
HOTAIRM1	HOX antisense intergenic RNA myeloid 1
HOX	Homeobox
HOXA9	Homeobox A9
HSCs	Hematopoietic Stem Cells
HSPCs	Hematopoietic Stem and Progenitor Cells
HTS	High-Throughput Screen
IDH	Isocitrate dehydrogenase
IDH	Isocitrate Dehydrogenase
ITD	Internal Tandem Duplication
JARID1	Jumonji AT-Rich Interactive Domain 1
KLF1	Kruppel-Like Factor 1
LICs	Leukemia Initiating Cells

LMO2	LIM Domain Only 2
LMPP	Lymphoid-primed Multipotent Progenitors
lncRNA	long non-coding RNA
LSCs	Leukemic Stem Cells
LSD1	Lysine-Specific Demethylase 1
MAPK	Mitogen-Activated Protein Kinase
MDS	Myelodysplastic Syndromes
MDS/MPN	Myelodysplastic/Myeloproliferative Neoplasm
MEIS1	Myeloid Ecotropic Viral Integration Site 1
MFC	Multiparameter Flow Cytometry
MGF	Mas cell Growth Factor
MHC	Major Histocompatibility Complex
miRISC	miRNA-induced silencing complex
miRNA	microRNA
MLL	Mixed Lineage Leukemia
MPAL	Myeloid or mixed Phenotype Acute Leukemia
MPP	Multipotent Progenitors
MRD	Minimal Residual Disease
MSCs	Mesenchymal Stem Cells
NAP1L3	Nucleosome Assembly Protein 1 Like 3
NGS	Next Generation Sequencing
NHEJ	Non-homologous End Joining
NK	Natural killer
NPM1	Nucleolar Phosphoprotein Member 1
NSG-SGM3	NOD Scid IL2Rgnull-3-SCF/GM/IL3
Ocn	Osteocalcin
Osx	Osterix



PAM	Protospacer Adjacent Motif
PBX3	Pre-B-Cell Leukemia Homeobox 3
PcG	Polycomb-Group
PCR	Polymerase Chain Reaction
PHF6	PHD Finger Protein 6
PI	Propidium Iodide
piRNA	piwi-interacting RNA
PRC1/2	Polycomb Repressive Complex 1/2
pre-GM	pre-Granulocyte-Macrophage progenitors
PTD	Partial Tandem Duplications
PTH	Parathyroid Hormone
PU.1-AS	PU.1 antisense transcript
RB1	Retinoblastoma 1
RBC	Red Blood Cells
RFP	Red Fluorescent Protein
RING1/2	Really Interesting New Gene 1/2
RISC	RNA induced silencing complex
RNAi	RNA interference
rRNA	ribosomal RNA
RT-qPCR	Real-Time quantitative PCR
s-AML	secondary AML
SAMHD1	SAM and HD Domain 1
SBDS	Shwachman-Bodian-Diamond Syndrome
Sc	Scramble
scaRNA	small cajal body-specific RNA
SCF	Stem Cell Factor
SF3B1	Splicing Factor 3b Subunit 1

sgRNA	single guide RNA
SHC	Src Homology 2 Domain Containing
shRNA	short hairpin RNA
SIPA1	Signal-Induced Proliferation-Associated gene 1
SIV	Simian Immunodeficiency Virus (SIV)
snoRNA	small nucleolar RNA
snRNA	small nuclear RNA
SRSF2	Serine and Arginine rich Splicing Factor 2
STAT5a	Signal Transducer and Activator of Transcription 5A
Suz12	Suppressor of Zeste 12
t-AML	therapy-related AML
TAL1	T-Cell Acute Lymphocytic Leukemia 1
TdT	Terminal deoxynucleotidyl Transferase
TET2	Tet methylcytosine dioxygenase 2
TFs	Transcription Factors
TK	Thymidine Kinase
TKD	Tyrosine Kinase Domain
TP53	Tumor Protein P53
TPO	Thrombopoietin
tRNA	transfer RNA
U2AF1	U2 Small Nuclear RNA Auxiliary Factor 1
UCBs	Umbilical Cord Blood Cells
VCAM-1	Vascular Cell Adhesion Molecule 1
VLPs	Virus-Like Particles
WBCs	White Blood Cells
WHO	World Health Organization

# **1 INTRODUCTION**

## **1.1 HEMATOPOIETIC SYSTEM**

The hematopoietic system includes the bone marrow (BM), spleen, lymph nodes and thymus, which are involved in the production of various blood cell types within the body. Specialized blood cells perform different functions in the body including oxygen supply, contributing to wound healing and protection from pathogens.

Oxygen is a vital component of all human cells, essential for cell growth and energy production. Red blood cells (RBCs) are responsible for oxygen transfer from the lung to all cells in the human body. Oxygen transport from the lung to the blood and uptake by cells is performed by passive diffusion (Krogh 1919). The protein hemoglobin in RBCs facilitates the transferring of 98% of oxygen in blood via reversible binding, whilst the rest of oxygen is found in free form in plasma and inside erythrocytes (Popel 1989).

Blood cells play an important role during hemostasis to stop bleeding, induce inflammation and prevent infection. In the wound healing process, numerous types of blood cells including platelets, neutrophils, monocytes, lymphocytes and dendritic cells are involved. The wound repair process is divided into three stages: inflammation, new tissue formation and remodeling (Gurtner, Werner et al. 2008). In the first stage, blood cells (immune cells and platelets) activate the coagulation cascade (hemostasis), inflammation pathways and prime anti-infection machinery. Platelets aggregate at the wound site and induce plug formation, after which insoluble fibrin forms a mesh to strengthen and stabilize the blood clot (Gale 2011).

Blood cells also contribute to both types of immunity, the innate and adaptive responses. The major blood cells and their products in the innate immune system include monocytes, macrophages, neutrophils and natural killer (NK) cells. The central players in the innate immune system are neutrophils, which are recruited and activated at the site of infection to eliminate pathogens (Witko-Sarsat, Rieu et al. 2000). NK cells are morphologically similar to B and T lymphocytes, but unlike lymphocytes they do not have specific antigen receptors. NK cells recognize abnormal cells through either FcR (immunoglobulin receptors) or their receptors for MHC (major histocompatibility complex) class I (Parkin and Cohen 2001). The adaptive immune system consists of two major types of lymphocytes: T cell are involved in cell-mediated response and B cells are mainly responsible for antibodies-related (humoral) immunity.

### 1.1.1 Hematopoiesis

Hematopoiesis describes the procedure of formation, development and differentiation of blood cells. In a healthy adult human, the blood system continuously produces approximately  $10^{11}$ - $10^{12}$  blood cells daily (Beerman, Maloney et al. 2010, Doulatov, Notta et al. 2012, Harrison 1979). Here, the term of blood cells includes RBCs, platelets and white blood cells (WBCs). The main hematopoietic cell types and their functions are summarized in Table 1.

The most common cells found in blood are RBCs, also named erythrocytes. In the human body,  $2-3 \times 10^6$  RBCs are produced every second in the BM with a life span of 120 days. Macrophages in the spleen or liver remove old or damaged RBCs from the blood circulation. Platelets originating from megakaryocytes play a role in hemostasis, thrombosis, inflammation and the immune system with an average life time of 8-10 days (Ho-Tin-Noe 2018).

WBCs are divided into two major subgroups: lymphoid and myeloid cells. The lymphoid cells consist of T cells, B cells and NK cells, which have roles in innate and adaptive immunity. The myeloid group includes monocytes, granulocytes (neutrophils, eosinophils and basophils), megakaryocytes and erythrocytes.


In mammalian adults, all blood cell types originate from a rare cell type in the BM termed hematopoietic stem cells (HSCs).

**Table1. Hematopoietic cell types and their function.**

Progenitors	Subtype	Function
Myeloid cells	Eosinophils	Involved in hypersensitivity and helminth infection
	Granulocytes (neutrophils)	Ingestion and destruction of microorganism
	Basophil	Inflammatory reactions and acute and chronic allergic development
	Macrophages	Derived from monocytes, respond to foreign material and release substances to stimulate other immune cells
	Megakaryocytes	Give rise to platelets
	Monocytes	Circulate in blood and migrate to tissue, they differentiate into macrophages
	Erythrocytes	Oxygen delivering
	Mast cells	Involved in allergy, anaphylaxis, wound healing, angiogenesis
Lymphoid cells	Cytotoxic T cells	Eradicate virus-infected cells as well as tumor cells
	T helper ( $T_H$ ) cells	Produce cytokines to activate B cells and T cells
	Memory T cells	Previously encountered and respond to their cognate antigen and ready to respond faster and stronger to the same antigen
	B cells	Produce antibodies
	Dendritic cells	Process antigen and present to T cells
	Natural killer cells	Part of innate immune system to kill viral infected cells and cancer cells

### 1.1.2 Hematopoietic stem cells and differentiation

HSCs are characterized by a self-renewal capacity to generate more stem cells (clonal expansion) and their ability to differentiate into various kinds of progenitor cells (clonal extinction) (Pina and Enver 2007). Indeed, HSCs represent the top cellular hierarchy of blood cells, from which progenitors derive, that give rise to common precursor cells (Figure 1). Studies have shown that only approximately 1000 HSCs contribute to hematopoiesis and peripheral blood production (Catlin, Busque et al. 2011). Early *in vivo* HSC tracing experiments performed in mice, revealed that the average cell division time of a quiescent HSC is 57 days (Cheshier, Morrison et al. 1999). Interestingly, other studies showed that dormant HSCs divided every 145 days or five times in a mouse's lifespan (Wilson, Laurenti et al. 2008). Dormant HSCs (d-HSCs) are thought to only serve in injury situations.



**Figure 1. Hierarchy model of hematopoiesis.** Hematopoietic stem cells (HSCs) with self-renewal capacity differentiate into progenitor cells and all mature blood cells. Figure reprinted with permission from the publisher (Wikipedia 2018).


In humans, it is not feasible to estimate the division time of HSCs by BrdUrd or H2-B labelling, but by using telomere length or the changing ratio of maternal/parental X-chromosome with age, the HSC replication rate was estimated to be on average once every 40-45 weeks (Catlin, Busque et al. 2011, Shepherd, Gutterop et al. 2004). However, considering mouse and human lifespans, the division number of HSCs is roughly similar between mice and humans.

### Origin of HSCs

During embryogenesis, blood cells originate from the mesodermal layer. The first wave of blood cell development occurs in the yolk sac (day E7.5 in mouse) and is termed “primitive” or embryogenic hematopoiesis, as it produces primitive nucleated erythroid, macrophage and megakaryocytes progenitors (Figure 2) (Palis, Robertson et al. 1999, Tober, Koniski et al. 2007). The second hematopoietic wave in the yolk sac, embryo proper and allantois generates erythroid and myeloid progenitors (EMP); B-1a and T lymphocytes (Dzierzak and Bigas 2018). Thus, the formation of blood cells in the first and second wave occurs without existing of HSCs. In the mouse embryo, HSCs originate at day E9.5 from endothelial cells expressing vascular endothelial cadherin (Zovein, Hofmann et al. 2008).

**Figure 2. Development of hematopoiesis in mice.**

**A.** The position of hematopoiesis first in yolk sac (YS) blood island then in aorta-gonad mesonephros (AGM) at day 10.5 and later in fetal liver, spleen and BM. **B.** The blood lineage hierarchy in each step of hematopoiesis. ECs (endothelial cells), RBCs (red blood cells), LTHSC (long-term hematopoietic stem cell), ST-HSC (short-term hematopoietic stem cell), CMP (common myeloid progenitor), CLP (common lymphoid progenitor), MEP (megakaryocyte-erythroid progenitor), GMP (granulocyte-monocyte progenitor). **C.** time frame and location of hematopoiesis. Figure reprinted with permission from the publisher (Orkin and Zon 2008).



Hematopoiesis converts to “definitive” in aorta-gonad mesonephros (AGM), fetal liver and eventually in the BM (adult hematopoiesis) (Dzierzak and Medvinsky 1995, Galloway and Zon 2003).

## **HSC Heterogeneity**

For a long time, HSCs were thought to be one homogenous population which gave rise to multipotent progenitors (MPPs) which in turn differentiated into different lineages. However, recent findings demonstrate that HSCs are heterogenous. Indeed, single-cell transplantation of HSCs showed a broad range of variability in the reconstitution and self-renewal capacity of mouse HSCs (Dykstra, Kent et al. 2007, Morita, Ema et al. 2010, Yamamoto, Morita et al. 2013). Taking advantage of new *in vivo* imaging technology, HSCs are now classified as having long-, intermediate- and short-term self-renewal capacities based on their specific cell surface markers (Doulatov, Notta et al. 2012, Notta, Doulatov et al. 2011). Aside from their heterogeneity in self-renewal capacity, single-cell transplantation experiments also demonstrated that the majority of HSCs are biased towards certain lineages during differentiation and only a few can differentiate equivalently to produce all mature blood cells (Carrelha, Meng et al. 2018, Dykstra, Kent et al. 2007, Morita, Ema et al. 2010, Yamamoto, Morita et al. 2013). Additionally, single-cell RNA sequencing revealed the heterogeneity of RNA expression profiles amongst HSCs, which reflect their ultimate destination of differentiation (Adolfsson, Mansson et al. 2005, Velten, Haas et al. 2017). Thus, all current data suggest that transcriptional lineage programming determines HSC function and is associated with HSC lineage commitment.

Together, considering the limitations to isolate pure HSCs and using different definition and markers to identify HSCs in studies, HSCs are a heterogeneous population with varying RNA expression profiles, differing differentiation abilities and distinct reconstitution and self-renewal capacities.

## **Isolation of hematopoietic stem cells**

As discussed above, HSCs are very rare, which makes their deep characterization difficult. For example, in human BM, where the HSCs reside, only 1 in a million cells represents a HSC (Wang, Doedens et al. 1997). So far, no unique marker for HSCs has been discovered.

For practical reasons, HSCs have mainly been studied in the mouse. To immune phenotypic isolation of mouse HSCs from the BM, HSCs were first defined as Lin<sup>-</sup>cKit<sup>+</sup>Sca-1<sup>+</sup> (LSK)


cells in 1992 (Ikuta and Weissman 1992). Among the LSK cells, functional studies revealed that the CD34<sup>+</sup> population was significantly enriched for long-term HSCs (Osawa, Hanada et al. 1996). Later, the HSCs could be further enriched to constitute 35-50% of the isolated cells, using an alternative sorting protocol defined as the SLAM phenotype (CD150<sup>+</sup>CD48<sup>-</sup>) (Kiel, Yilmaz et al. 2005).

To isolate and study HSCs in humans, The surface marker CD34 was suggested to enrich HSCs as it is found on less than 5% of all blood cells (Civin, Strauss et al. 1984). Although more than 99% of all human HSCs are CD34<sup>+</sup>, several studies have shown the existence of CD34<sup>-</sup> HSCs (Ishii, Matsuoka et al. 2011). Further studies have shown that HSCs in humans can be further enriched in cells not expressing CD38 or CD45RA (Bhatia, Wang et al. 1997, Lansdorp, Sutherland et al. 1990). Another interesting marker is CD90 (Thy1) which has been used to obtain highly purified HSCs (Baum, Weissman et al. 1992). Therefore, Lin<sup>-</sup>CD34<sup>+</sup>CD38<sup>-</sup>CD90<sup>+</sup>CD45RA<sup>-</sup> is an immune phenotype that has been widely used to study and isolate human HSCs in the field, although this protocol is continuously being updated with additional markers. For example, including CD49f marker in Lin<sup>-</sup>CD34<sup>+</sup>CD38<sup>-</sup>CD90<sup>+</sup>CD45RA<sup>-</sup> cells led to further purification of long-term repopulating HSCs (Notta, Doulatov et al. 2011). Although the research field has not been able to isolate a completely pure population of HSCs, which might reflect their heterogeneity, we can now enrich them for further study of HSCs. A summary of specific markers used for isolation of each HSC and progenitor population in mouse and human is depicted in Figure 3.

## **Bone marrow niche**

The concept of a niche for hematopoietic stem cells was first suggested in 1978 (Schofield 1978). This hypothesis proposed that HSCs associated with other cells in BM. BM is a semi-solid tissue located in the spongy or cancellous portions of the tibia, femur, ribs, sternum, vertebrae and pelvis. The BM niche regulates various HSC activities including self-renewal, differentiation, mobilization, and engraftment. The main cell types in the BM niche are osteolineage cells, perivascular cells, endothelial cells, adipocytes, macrophages and mesenchymal stem cells (MSCs) (Figure 4).






**Figure 3. Cell surface markers of main classes of stem cells and progenitors in mouse and human.** Figure reprinted with permission from the publisher (Doulatov *et al.* 2012).

It has previously been reported that various types of osteolineage cells reside in the BM with distinct roles regulating different lymphoid cells. Knockout of osteocalcin (Ocn)-expressing osteolineage cells results in a loss of T cells (Yu, Saez *et al.* 2015), whereas osterix-expressing (Osx<sup>+</sup>) and Col(I)a2.3-expressing osteolineage cell deletion impairs the maturation of B cell progenitors (Visnjic, Kalajzic *et al.* 2004, Zhu, Garrett *et al.* 2007). Many HSCs localize adjacent to the blood vessels, which suggests that the perivascular region is crucial for HSC maintenance. In addition, MSCs also reside near vessels and can differentiate into osteolineage cells, chondrocytes, and adipocytes. Genetic deletion of MSCs in mice has recently been shown to affect the HSCs (Mendez-Ferrer, Battista *et al.* 2010), for example SIPA-1 deletion leads to the development of myelodysplastic/myeloproliferative neoplasm (MDS/MPN) phenotype in mice (Xiao, Dolinska *et al.* 2018). Another cell type which surrounds sinusoidal endothelial cells, or is located near to the endosteum, is CXC chemokine ligand (CXCL)12-abundant reticular (CAR) cells which co-localize with HSCs (Sugiyama,

Kohara et al. 2006). Ablation of CAR cells dramatically affects adipogenic and osteogenic differentiation, whilst decreased production of stem cell factor (SCF) and CXCL12 cytokines, subsequently caused a reduction in erythroid progenitors, cycling lymphoid cells and HSCs (Omatsu, Sugiyama et al. 2010).

Endothelial cells secrete specific growth factors in a paracrine-specific manner, known as angiocrine factors, and balance self-renewal and support expansion of hematopoietic stem and progenitor cells (HSPCs) over differentiation (Kobayashi, Butler et al. 2010). Moreover, they express adhesion molecules including P-selectin, E-selectin, vascular cell adhesion molecule 1 (VCAM-1), and intercellular adhesion molecule 1 (Mazo, Gutierrez-Ramos et al. 1998, Rafii, Mohle et al. 1997) which are important for HSCs homing into the BM. The percentage of adipocytes in the BM or fatty marrow increases with age in both humans and mice, yet HSC function decreases, implying that adipocytes negatively regulate HSCs (Yu and Scadden 2016).



**Figure 4. HSC niche in the adult bone marrow.** Schematic of various stromal cells types and extrinsic signals including growth factors, cytokines, morphogens, extracellular matrix proteins, and adhesion molecules in the adult BM that contribute to the maintenance and regulation of HSCs. Figure reprinted with permission from the publisher (Yu and Scadden 2016).

Macrophages are another key cell type in the BM niche. Depletion of macrophages caused loss of osteoblasts, significant reduction of HSC-trophic cytokines and HSC mobilization into the blood (Winkler, Sims et al. 2010). Interestingly, macrophages supported erythroid lineage development (Palis 2016).

New findings have suggested that HSCs reside in different spots within the BM and depending on their location may have different functions. For example, some studies have shown that osteoblastic cells regulate HSCs via N-cadherin (Calvi, Adams et al. 2003, Zhang, Niu et al. 2003). Yoshihara suggested HSCs close to thrombopoietin-producing osteoblasts remain in a quiescent state (Yoshihara, Arai et al. 2007), whilst another study revealed that the vascular cells regulate HSCs migration to the vascular niche via CXCL12 (Kiel and Morrison 2006). Altogether, it seems that HSCs localize to a specific area within the BM and are surrounded by hematopoietic as well as non-hematopoietic cells, which in turn provide a tonic regulation of HSCs. Therefore, any perturbation of these non-hematopoietic cells may cause abnormal function of HSCs.

### **1.1.3 Epigenetic regulation of hematopoiesis**

The term “epigenetic” (Epi is Greek prefix means “over”, “outside of”) was suggested by Conrad Waddington in 1942 (Waddington 2012). Epigenetics refers to the stable and heritable modifications in gene activity that occur without any alteration in DNA sequence, including histone modification, DNA methylation and non-coding RNAs. Epigenetic regulation is a complex process by combination and interaction of many epigenetic modifiers which add “writers” or remove “erasers” modifications on histones or DNA and recognize and respond to the modifications “readers”.

In eukaryotes DNA is wrapped around a histone octamer called the nucleosome and nucleosome units fold into chromatin structure. Epigenetic regulatory factors modify chromatin structure in both a global and gene-specific manner by post-translational modification of histones and/or DNA methylation to provide or prevent access of transcription factors (TFs) to promoter regions. In other words, epigenetic modifications can result in either “loose” chromatin regions, called euchromatin, which are available for TFs or “tightly” packed chromatin regions, termed heterochromatin, which are inaccessible for transcription. For example, transcriptionally silenced regions in eukaryotic cells are associated with methylation of DNA in CpG regions and histone H3 dimethylated on lysine 9 (H3K9me2) and histone H3 trimethylated on lysine 27 (H3K27me3) whereas transcriptionally active regions contain high levels of H4 trimethylated on lysine 4

(H3K4me3) and lack DNA methylation (Li, Carey et al. 2007). Transcription of a gene occurs on naked DNA by TFs and RNA polymerase enzymes. Therefore, epigenetic mechanisms to alter chromatin and provide naked DNA for co-activators and TFs is the key step to determine cell identities.

The epigenetic landscape was first suggested by Waddington in 1957 to explain the role of epigenetic factors to canalize cells, originating from stem cells, during differentiation (Ferrell 2012). Therefore, epigenetic factors stabilize gene expression profile in cells and canalize cell-type identities. Transcription and epigenetic regulatory factors play crucial roles in keeping the balance between HSC maintenance and differentiation. However, differentiation in the hematopoietic system is not as simple as Waddington's original theory. First, in the beginning of hematopoiesis (the first and second wave) in the embryo stage, differentiated cells are made prior to and independently of HSCs. Furthermore, it was confirmed that by inducing expression of specific lineage TFs in hematopoietic cells, it is possible to switch the cells from one lineage to another (Graf 2002). Below, we will discuss the important epigenetic factors which play a role in hematopoiesis.

## **DNA methylation**

DNA methylation was first suggested in 1975 as an epigenetic mechanism of imprinting to inactivate the X-chromosome in female cells (Holliday and Pugh 1975, Riggs 1975). In human cells, DNA methylation predominantly occurs in cytosine at the C5 position. DNA methylation mainly occurs in CpG dinucleotides (60-80% of all CpGs are methylated) at the promoter regions which influences transcriptional activity (Lister, Pelizzola et al. 2009, Saxonov, Berg et al. 2006).

Stem cell differentiation is linked to gradual methylation of CpG islands by DNA methyltransferases (Trowbridge and Orkin 2011). In hematopoiesis, different methylation patterns are significantly correlated with the global gene expression pattern and results in cell fate decisions (Ji, Ehrlich et al. 2010). DNA methylation might play a role as an epigenetic gatekeeper to retain a specific-cell lineage pattern during differentiation.

Two methyltransferases in particular, DNMT3a and DNMT3b are required for *de novo* methylation. During differentiation of HSCs, DNMT3a and DNMT3b are involved in the silencing of genes which regulate HSC self-renewal activity (Trowbridge and Orkin 2011). Hence, deletion of *DNMT3a* and *DNMT3b* in HSCs caused upregulation of self-renewal genes and subsequently increased HSC self-renewal and impaired HSC differentiation

(Challen, Sun et al. 2014). Another methyltransferase, DNMT1 preserves the DNA methylation patterns during DNA replication. DNMT1 also has an important role during HSC differentiation towards multipotent progenitors of a myeloid-restricted lineage (Trowbridge, Snow et al. 2009). Genes that are important in maintaining HSCs, including *Meis1*, *Hoxa9* and *Prdm16*, are heavily methylated and so transcriptionally silenced in progenitor cells during differentiation (Kosan and Godmann 2016). Whereas, active genes in differentiated cells are found to be methylated in HSCs and selectively become demethylated during the lineage commitment process. For example, *Gadd45a* a critical gene in myeloid development was found to be upregulated and demethylated in cells transitioning from common myeloid progenitor (CMP) to granulocyte-monocyte progenitor (GMP) stage whereas it is methylated and silenced in HSCs (Ji, Ehrlich et al. 2010).

## **Histone modifications**

There are many post-translational histone modifications which influence chromatin structure including acetylation, methylation, phosphorylation, ubiquitylation, sumoylation, adenosine diphosphate (ADP) ribosylation, and deamination (Kouzarides 2007). These modifications are associated with various processes including transcription regulation, DNA repair, DNA replication and condensation (Kouzarides 2007). Histone-modifying enzymes are summarized in Table 2.

There are some important histone-modifying enzymes in hematopoiesis which play key roles in HSC function including self-renewal and differentiation. Important histone acetyltransferase (HAT) that transfer acetyl groups to specific lysine residue of histones in hematopoiesis are p300/CBP (CBP and p300), MYST (Tip60, MOZ, MORF, HBO1 and HMOF), and GNAT (PCAF, Gnc5 and ELP3) (Sun, Man et al. 2015). P300 is associated with promoting differentiation of HSCs by acetylation of the *C-Myb* promoter region, whereas CBP acetylates *Gfi1b* and promotes self-renewing of HSCs and blocks differentiation (Sun, Man et al. 2015).

Polycomb-group (PcG) proteins are histone modifiers which exist in two main complexes; polycomb repressive complex 1 (PRC1) and 2 (PRC2). They have been shown to be involved in gene repression in HSCs during self-renewal and differentiation (Radulovic, de Haan et al. 2013).

**Table 2. Histone-modifying enzymes.** Table adapted with permission from the publisher (Kouzarides 2007).

Function	Name	Residues modified
Acetyltransferase	HAT1	H4 (K5, K12)
	CBP/P300	H3 (K14, K18), H4 (K5, K8), H2A (K5), H2B (K12, K15)
	PCAF/GCN5	H3 (K9, K14, K18)
	TIP60	H4 (K5, K8, K12, K16), H3K14
	HB01	H4 (K5, K8, K12)
Deacetylase	SirT2	H4K16
Lysine Methyltransferase	SUV39H (1-2)	H3K9
	GLP/EHMT1	H3K9
	G9a/EHMT2	H3K9
	ESET	H3K9
	CLL8	H3K9
	MLL (1-5)	H3K4
	SET1 (A-B)	H3K4
	ASH1	H3K4
	SET2	H3K36
	NSD1	H3K36
	SYMD2	H3K36
	DOT1	H3K79
	Pr-SET 7/8	H4K20
	SUV4 20H(1-2)	H4K20
	EZH2	H3K27
	RIZ1	H3K9
Lysine Demethylase	LSD1/BHC110	H3K4
	JHDM1 (a-b)	H3K36
	JHDM2 (a-b)	H3K9
	JMJD2A/JHDM3A	H3 (K9, K36)
	JMJD2B	H3K9
	JMJD2C/GASC1	H3 (K9, K36)
	JMJD2D	H3K9
Arginine Methyltransferase	CARM1	H3 (R2, R17, R26)
	PRMT4	H4R3
	PRMT5	H3R8, H4R3
Serine/Threonine Kinase	Haspin	H3T3
	MSK (1-2)	H3S28
	CKII	H4S1
	Mst1	H2BS14
Ubiquitilases	Bmi/Ring1A	H2AK119
	RNF20/RNF40	H2BK120
Proline Isomerase	ScFPR4	H3 (P30, P38)

PRC2 includes the enhancer of zeste (E(z)), Suppressor of zeste 12 (Suz12) and Extra sex combs (Esc). Overall, PRC2 is responsible for the methylation of H3K27 (H3K27me2 and H3K27me3) at the promoter sites which is associated with transcriptional repression of target genes. PRC1 consists of BMI1, RING1/2, MEL-18, RAE28/MPH1, and M33/CBX2. The PRC1 complex binds to H3K27me3, which is established by PRC2, and stabilizes gene silencing by H2A ubiquitination (H2AK119ub1). Ubiquitination of H2A as the last step of

gene repression is thought to inhibit RNA pol II in the initiation or elongation phase of transcription (Radulovic, de Haan et al. 2013). Deletion of *Bmi1*, part of PRC1, in mice caused a severe reduction in the number of long-term HSCs (Park, Qian et al. 2003), whilst deletion of *Mel-18* induced self-renewing of HSCs and a defect in B cells, suggesting a role for PRC1 in self-renewal and differentiation of HSCs (Akasaka, Tsuji et al. 1997, Kajiume, Ninomiya et al. 2004).

Mixed Lineage Leukemia (MLL), lysine methyl transferase enzyme, plays an important role in the second wave of embryo hematopoiesis, AGM and further HSC development (Orkin 2000). MLL is associated with transcriptional activation by trimethylation of H3 (H3K4me3) at downstream target genes, particularly *HOX* genes.

Two main members of the histone demethylase family are Lysine-specific demethylase 1 (LSD1) and the Jumonji C (JmjC) domain-containing family. LSD1 is part of a complex with histone deacetylases (HDACs) and CoREST and acts as a transcriptional repressor to downregulate expression of target genes by removing one or two methyl groups from H3K4 or H3K9. Several JmjC domain proteins have been found including JHDM1, JHDM2, JMJD2 and Jumonji AT-rich interactive domain 1 (JARID1). Jarid1b is highly expressed in HSCs and is important for HSC self-renewal (Stewart, Albert et al. 2015).

Some histone methylations such as H3K4, H3K36, and H3K79 are linked with gene activation, whereas methylation on H3K9, H3K27, and H4K20 results in transcription inhibition. Interestingly, many promoters in stem cells co-exist alongside transcriptional activation (H3K4me3) and repression marks (H3K27me3). These bivalent domains allow stem cells to rapidly activate or silence genes during differentiation (Voigt, Tee et al. 2013).

## **Non-coding RNA**

RNAs that do not translate to proteins are defined as non-coding RNAs, which include ribosomal RNA (rRNA), transfer RNA (tRNA), small nuclear RNA (snRNA), small nucleolar RNA (snoRNA), small cajal body-specific RNA (scaRNA), piwi-interacting RNA (piRNA), microRNA (miRNA) and long non-coding RNA (lncRNA). Some non-coding RNAs, in particular miRNA and lncRNA, play a role in gene regulation and are therefore regarded as part of the epigenetic regulators.

miRNAs are short RNAs containing 21-24 bases and are responsible for post-transcriptional gene silencing by pairing to the complement region of target mRNA and recruiting miRNA-induced silencing complexes (miRISC) (Fabian and Sonenberg 2012). Several reports have

shown that miRNAs play a pivotal function in hematopoiesis. In 2004, miRNA-181 was found to play a role as a regulatory factor in B cell differentiation (Chen, Li et al. 2004). In addition, growth factor independent-1 (Gfi1); a critical transcription factor for granulocytic differentiation is regulated by miR-21 (Velu, Baktula et al. 2009). In erythroid cell differentiation miR-92a and miR-17 targets regulatory erythroid genes (Li, Vecchiarelli-Federico et al. 2012). miRNAs are not only important during differentiation but are also crucial in the regulation of HSPC expansion, for example overexpression of miR-17-92 induced proliferation of multipotent hematopoietic progenitors (Li, Vecchiarelli-Federico et al. 2012).


lncRNAs contain more than 200 bases and it is estimated that 10,000-60,000 lncRNAs are expressed in human cells (Wilkes, Repellin et al. 2017). lncRNAs regulate gene expression at various steps including mRNA processing, during translation or protein synthesis, modification of mRNA stability, miRNA inhibition and as a scaffold to facilitate mRNA synthesis (Geisler and Collier 2013). Numerous studies have linked lncRNAs to hematopoiesis. For example, over 100 lncRNAs are exclusively expressed during erythroid development, some of which target the transcription factors *GATA*, T cell acute lymphocytic leukemia protein 1 (*TALI*) or Kruppel-like factor 1 (*KLF1*) (Geisler and Collier 2013). During myeloid development, PU.1 antisense transcript (PU.1-AS) was previously identified to target *PU.1* mRNA and downregulates *PU.1* transcription factor (Ebralidze, Guibal et al. 2008). Similarly, HOX antisense intergenic RNA myeloid 1 (HOTAIRM1) was identified as a myeloid-specific lncRNA which is highly expressed during granulocytic differentiation, whilst HOTAIRM1 suppresses *HOXA1* and *HOXA4* during myeloid differentiation (Zhang, Lian et al. 2009). Interestingly, more than 3000 lncRNAs have been found to be expressed during lymphoid development, suggesting that it is possible to characterize different lymphoid committed cells by the lncRNA expression patterns (Casero, Sandoval et al. 2015). Altogether, these studies indicate the important role of non-coding RNAs as part of the epigenetic regulatory compartment in HSC function and during differentiation of progenitor cells.

#### **1.1.4 Transcription factors involved in hematopoiesis**

Another intrinsic element which determines the identity of cells is the particular TF network to induce expression of lineage-specific genes. Critical TFs in hematopoiesis, contain nearly all groups of DNA-binding proteins rather than favoring a specific class. Some TFs are required for the development and maintenance of HSCs, some for the differentiation process



and some are involved in both. For example *MLL*, *Runx1*, *TEL/ETV6*, *SCL/tal1* and *LMO2* are “HSCs TFs” but *PU.1*, *Gfi-1*, *C/EBP $\alpha$*  are more lineage-restricted factors (Orkin and Zon 2008). A summary of some important TFs in hematopoiesis is depicted in Figure 5.



**Figure 5. Important transcription factors in hematopoiesis.** Important TFs in each stage depicted in red. The TFs in light font have not yet been identified translocated or mutated in human/mouse hematologic malignancies but the rest are associated with hematopoietic malignancies. LT-HSC (long-term hematopoietic stem cell), ST-HSC (short term hematopoietic stem cell), CMP (common myeloid progenitor), CLP (common lymphoid progenitor), MEP (megakaryocyte/erythroid progenitor), GMP (granulocyte/macrophage progenitor) RBCs (red blood cells). Figure reprinted with permission from the publisher (Orkin and Zon 2008).

## Factors required for development and maintenance of HSCs

Development and maintenance of HSCs is pre-determined by a series of TFs, which are influenced by the microenvironment, signaling pathways and epigenetic regulators. During the emergence of HSCs, signaling molecules originating from the adjacent germ cell layer induce expression of crucial TFs required for this step. Therefore, physical interaction between primitive endoderm and adjacent mesoderm and subsequently endodermal signaling are required for the development of hematopoietic cells during the embryo stage (Belaoussoff, Farrington et al. 1998). Some important TFs required for programming the ventral mesoderm toward HSCs are basic-helix-loop-helix (bHLH), SCL/TAL-1 and LMO2. The absence of these factors leads to defects in both endothelial and hematopoietic cells (Kim and Bresnick 2007). During the yolk stage, they are also required for development of blood cells, therefore they are pivotal for the development of primitive and definitive HSCs.

One of the key TFs for definitive hematopoiesis is Runx1, which is expressed during the first wave of hematopoiesis in mesoderm (North, Gu et al. 1999). Zebrafish models have shown that *Runx1*, *Lmo2* and *Scl* are regulated by Notch signaling, which induce expansion of HSCs (Burns, Traver et al. 2005). In adult human blood cells, Runx1 is expressed in all blood cells except erythrocytes (North, Stacy et al. 2004). *Runx1* and *Runx3* double-knockout caused BM failure and myeloproliferative disorder in mice via the non-transcriptional function of *Runx* in DNA repair (Wang, Krishnan et al. 2014). Apart from Runx1, another important factor for fetal HSCs is Sox17, which is uniquely expressed in HSCs during embryogenesis. Deletion of *Sox17* results in a lack of adult HSCs (Kim, Saunders et al. 2007).

Other TFs involved in hematopoiesis are *HOX* genes which classify in four clusters: *HOXA*, *HOXB*, *HOXC* and *HOXD*. HOX TFs are homeodomain-containing TFs which were originally characterized in *Drosophila*, yet with important roles in hematopoiesis (Shah and Sukumar 2010). They are expressed in HSCs and progenitors with specific expression profile patterns, dependent upon the cell lineage. For instance *HOXB3*, *HOXB4* and *HOXA9* are significantly expressed in uncommitted hematopoietic cells, while *HOXB8* and *HOXA10* are activated in myeloid-committed cells (Alharbi, Pettengell et al. 2013). Overexpression of *HOX* genes most often causes HSPC expansion and blockade of differentiation. Previously, *Hoxb6* overexpression has been demonstrated to increase the number of HSCs and myeloid progenitors whilst blocking erythropoiesis and lymphopoiesis (Fischbach, Rozenfeld et al. 2005).

The functional mechanism of *HOX* genes to regulate hematopoiesis is not yet clearly defined. However, some studies have suggested downstream targets of HOX TFs which are crucial in

hematopoiesis. For example, *HOXA9* activates expression of other *HOX* genes (*HOXA7* and *HOXA10*), *PBX3*, *MEIS1*, *Kit*, *Flt3* and *Sox4* which are important in hematopoiesis. Other important downstream target genes which are regulated by *HOX* genes are *GATA1*, *C-MYC* and *RUNX2* (Alharbi, Pettengell et al. 2013).

As mentioned above, one important upstream regulator of *HOX* is MLL, which directly regulates transcription of *HOX* genes. Another group of factors which directly regulate *HOX* genes are *CXD* genes (*CDX1*, *CDX2* and *CDX4*) (Brooke-Bisschop, Savory et al. 2017). Deletion of *Cdx1* and *Cdx2* in mice was shown to impair primitive hematopoiesis and resulted in a lack of primitive erythrocytes as well as an abnormal yolk sac vasculature (Brooke-Bisschop, Savory et al. 2017). Therefore, the *CDX* genes are also required for normal HSC function.

### **Important TFs for HSCs differentiation**

Two crucial factors which are not only important for HSC generation, but also important for differentiation of HSCs are GATA-2 and Runx-1. Inducible deletion of *Runx-1* in BM cells showed that *Runx-1* is essential for development of megakaryocytes and T and B cell differentiation, but is not essential for normal myeloid development (Ichikawa, Asai et al. 2004). Other studies, however, show that *Gata2* contributes to the generation and long-term maintenance of HSCs (de Pater, Kaimakis et al. 2013). Several pathways upstream of *Gata2* and *Runx1* regulate their expression at different stages of the HSC development and differentiation. The important regulatory pathways for GATA2 and RUNX1 are NOTCH (Guiu, Shimizu et al. 2013), BMP4 (Walmsley, Ciau-Uitz et al. 2002) and ETS/EGR (Taoudi, Bee et al. 2011). There are also some myeloid factors which are expressed in GMPs such as C/EBP $\alpha$  (Orkin and Zon 2008).

Another member of the GATA family which plays a crucial role in hematopoiesis is GATA1. Expression of *GATA1* is important for erythropoiesis (Fujiwara, Browne et al. 1996). Specific deletion of *Gata1* in megakaryocytes reduced platelet numbers and impaired the differentiation of megakaryocytes, which in turn increased numbers of immature megakaryocytes in the BM and spleen (Kuhl, Atzberger et al. 2005, Shivdasani, Fujiwara et al. 1997). Additionally, *GATA1* has also been shown to have an important function in the development of mast cells, eosinophils, basophils, and dendritic cells (Crispino and Horwitz 2017).

Activation of lineage-specific TFs determine differentiation of HSCs and usually the activated TFs in turn act as suppressors for other lineage-specific TFs (Orkin and Zon 2008). For instance, GATA-1 and PU.1 physically interact and antagonize the function of each other (Rhodes, Hagen et al. 2005). Expression of FOG in multipotent cells suppresses the eosinophil factor C/EBPbeta and yet during the eosinophil lineage commitment, C/EBPbeta blocks expression of FOG (Querfurth, Schuster et al. 2000). The TFs FLI-1 and EKLF cross-antagonise their respective functions during megakaryocyte or erythroid development (Starck, Cohet et al. 2003). Moreover, GFI-1 physically interacts with and suppresses PU.1, inducing macrophage differentiation (Dahl, Iyer et al. 2007). Interestingly, in HSC differentiation, during the phenomenon called “the GATA switch”, GATA2 activates GATA1 but after activation, GATA1 negatively regulates the expression of GATA2 (Grass, Boyer et al. 2003).

Altogether, the data support the fundamental role of TFs in hematopoiesis and Interestingly, somatic mutation or chromosomal translocations of the majority of TFs within the hematopoietic system are associated with hematopoietic malignancies.

## **1.2 ACUTE MYELOID LEUKEMIA**

Acute myeloid leukemia (AML) is a heterogeneous hematological malignancy which is characterized by clonal expansion of abnormal or immature myeloid progenitors in the BM and peripheral blood (O'Donnell, Abboud et al. 2012).

AML is the most common subtype of acute leukemia in adults, covering around 80% of all leukemia cases in adults (Yamamoto and Goodman 2008). The AML incidence rate is 1.3 per 100,000 in the population younger than 65 years old and increases with age to 12.2 patients per 100,000 in the population over 65 years old in the USA (De Kouchkovsky and Abdul-Hay 2016). Despite the development of therapeutic drugs and improvement of supportive care over the past 50 years, AML is only curative in 35-40% of adult patients under 60 years of age and 5-15% in patients over 60 years old (Dohner, Weisdorf et al. 2015). Therefore, there is a great demand for more efficient therapeutic strategies in AML.

### **1.2.1 Diagnosis of AML**

The general symptoms of AML patients result from an accumulation of poorly differentiated myeloid cells in the BM and peripheral blood leading to BM failure, anemia, thrombocytopenia, fatigue, anorexia, fever, night sweats and weight loss. Many symptoms

of AML are as a consequence of normal blood deficiency due to leukemia cells outcompeting normal cells in the BM. Therefore, patients do not have enough RBCs, WBCs and platelets.

Preliminary diagnosis of AML is based on morphology, notably the presence of 20% or more blast cells in at least 200 counted leukocytes on blood smear and 500 on speculated BM smear, with the exception of AML with t(15;17), t(8;21), inv(16), or t(16;16). (Dohner, Estey et al. 2017, Dohner, Estey et al. 2010). The nuclei of AML blasts are large in size and typically several nucleoli can be found in one AML cell.

Following morphology, subsequent diagnostic methods for AML are based on surface phenotyping. AML cells express CD markers that exist on normal myeloid cells, such as CD13, CD33 and CD34 (Campos, Guyotat et al. 1989). However, other CD markers might be expressed on AML cells dependent upon the AML subtype or stage of AML differentiation (Dohner, Estey et al. 2017). For example, AML cells may express CD4, CD14, CD11b, CD36 and CD64 (monocyte markers), CD36, CD71 and CD235a (erythroid markers), CD41 and CD61 (megakaryocyte markers). Moreover, AML cells may also express specific T or B cells markers; CD7, CD19, terminal deoxynucleotidyl transferase (TdT) and human leukocyte antigen-antigen D related (HLA-DR), (Dohner, Estey et al. 2017).

Finally, conventional cytogenetic analysis of AML cells is the next mandatory method for diagnosis and classification of AML cells which is based on chromosomal abnormalities. Fluorescence in situ hybridization (FISH) and new sequencing technologies are used to detect fusion genes (McKerrell, Moreno et al. 2016), following which molecular genetic screening for known mutations such as *NPM1*, *CEBPA*, *RUNX1*, *FLT3*, *TP53* and *ASXL1* should be done. The summary of tests or procedures for accurate diagnosis of AML is provided in Table 3 (Dohner, Estey et al. 2017).

**Table 3. Summary of tests and procedures to diagnose AML.**

<b>Cellular assays</b>
Blood count
Bone marrow aspirate
Immunophenotyping
<b>Genetic analysis</b>
Cytogenetics
Screening for mutations: <i>NPM1</i> , <i>CEBPA</i> , <i>RUNX1</i> , <i>FLT3</i> , <i>TP53</i> , <i>ASXL1</i>
Screening for fusion genes: <i>PML-RARA</i> , <i>CBFB-MYH11</i> , <i>RUNX1-RUNX1T1</i> , <i>BCR-ABL1</i> , <i>MLL-X</i>
<b>Additional procedures</b>
Medical history including exposure to toxic agents, prior malignancy and therapy, smoking
Family history to check germ line predisposition of myeloid neoplasms

### 1.2.2 Classification of AML

Several different classification systems have been suggested over the years based on morphologic analysis of blood smears and BM samples, expression levels of cell-surface or cytoplasmic markers using flow cytometry, cytogenetic karyotype tests and screening of genes that are important for AML development and maintenance.

As of 1970, AML was classified into eight major subtypes (M0-M7) according to the French-American-British (FAB) system, which is based on the morphology and immune phenotype of AML cells (Bennett, Catovsky et al. 1976). More recently, AML has been classified according to the world health organization (WHO) classification in 2008, which is based on leukemia-associated chromosomal translocations and inversions (Sabattini, Bacci et al. 2010). According to the WHO classification, AML can be divided in seven subtypes: (1) AML with chromosomal abnormalities, (2) AML with myelodysplasia-related changes, (3) therapy-related myeloid neoplasms, (4) AML not otherwise specified (NOS) (similar to FAB Classification M0–M7 with others such as acute megakaryoblastic leukemia (AMKL), acute panmyelosis (APMF) with myelofibrosis, and pure erythroleukemia), (5) myeloid sarcoma, (6) myeloid proliferations related to Down syndrome, and (7) blastic plasmacytoid dendritic cell neoplasm (Vardiman, Thiele et al. 2009). The most recent WHO classification was updated in 2016 (Table 4) (Arber, Orazi et al. 2016). In the updated version, a new class “myeloid neoplasms with germ line predisposition” was added (Table 5); for example, germline *CEBPA* mutations frequently detected in AML patients who need special genetic counselling for their families (DiNardo, Bannon et al. 2016, Pabst, Eyholzer et al. 2008).

Presently, genetic analyses of AML samples by next generation sequencing (NGS) has provided further information and better views into AML classification (Grimwade, Ivey et al. 2016, Papaemmanuil, Gerstung et al. 2016).

AML can also be classified based on etiology into three categories: (1) de novo AML, (2) therapy-related AML (t-AML), and (3) secondary AML (s-AML) originated from MDS (myelodysplastic syndromes) or other myeloid proliferative disorder (Lindsley, Mar et al. 2015).

**Table 4. WHO classification of AML and related precursor neoplasms and acute leukemia.**  
Table adapted with permission from the publisher (Dohner, Estey et al. 2017).

<b>AML with recurrent neoplasms</b>
<ul style="list-style-type: none"> <li>• AML with t(8;21)(q22;q22.1); RUNX1-RUNX1T1</li> <li>• AML with inv(16)(p13.1q22) or t(16;16)(p13.1;q22); CBFB-MYH11</li> <li>• Acute promyelocytic leukemia with PML-RARA<sup>1</sup></li> <li>• AML with t(9;11)(p21.3;q23.3); MLLT3-KMT2A<sup>2</sup></li> <li>• AML with t(6;9)(p23;q34.1); DEK-NUP214</li> <li>• AML with inv(3)(q21.3q26.2) or t(3;3)(q21.3;q26.2); GATA2,MECOM(EVI1)</li> <li>• AML (megakaryoblastic) with t(1;22)(p13.3;q13.3); RBM15-MKL1<sup>3</sup></li> <li>• Provisional entity: AML with BCR-ABL1</li> <li>• AML with mutated NPM1<sup>4</sup></li> <li>• AML with biallelic mutations of CEBPA<sup>4</sup></li> <li>• Provisional entity: AML with mutated RUNX1</li> </ul>
<b>AML with myelodysplasia-related changes<sup>5</sup></b>
<b>Therapy-related myeloid neoplasms<sup>6</sup></b>
<b>AML, NOS</b>
<ul style="list-style-type: none"> <li>• AML with minimal differentiation</li> <li>• AML without maturation</li> <li>• AML with maturation</li> <li>• Acute myelomonocytic leukemia</li> <li>• Acute monoblastic/monocytic leukemia</li> <li>• Pure erythroid leukemia<sup>7</sup></li> <li>• Acute megakaryoblastic leukemia</li> <li>• Acute basophilic leukemia</li> <li>• Acute panmyelosis with myelofibrosis</li> </ul>
<b>Myeloid sarcoma</b>
<b>Myeloid proliferations related to Down syndrome</b>
<ul style="list-style-type: none"> <li>• Transient abnormal myelopoiesis</li> <li>• Myeloid leukemia associated with Down syndrome</li> </ul>
<b>Blastic plasmacytoid dendritic cell neoplasm</b>
<b>Acute leukemias of ambiguous lineage</b>
<ul style="list-style-type: none"> <li>• Acute undifferentiated leukemia</li> <li>• MPAL with t(9;22)(q34.1;q11.2); BCR-ABL1<sup>8</sup></li> <li>• MPAL with t(v;11q23.3); KMT2A rearranged</li> <li>• MPAL, B/myeloid, NOS</li> <li>• MPAL, T/myeloid, NOS</li> </ul>

MPAL, mixed phenotype acute leukemia; NK, natural killer.

1. Other recurring translocations involving RARA should be reported accordingly: AML with t(11;17)(q23;q12); ZBTB16-RARA, t(11;17)(q13;q12); NUMA1-RARA, t(5;17)(q35;q12); NPM1-RARA, or AML with STAT5B-RARA.

2. Other translocations involving KMT2A (MLL) should be reported accordingly: AML with t(6;11)(q27;q23.3); MLLT4-KMT2A, AML with t(11;19)(q23.3;p13.3); KMT2A-MLLT1, t(11;19)(q23.3;p13.1); KMT2A-ELL, t(10;11)(p12;q23.3); MLLT10-KMT2A.

3. Rare leukemia most commonly occurring in infants.

4. Diagnosis is made irrespective of the presence or absence of multilineage dysplasia.

5. At least 20% blood or marrow blasts AND any of the following: previous history of MDS or MDS/MPN; myelodysplasia-related cytogenetic abnormality; multilineage dysplasia; AND absence of both prior cytotoxic therapy for unrelated disease and aforementioned recurring genetic abnormalities. Cytogenetic abnormalities sufficient to diagnose AML with myelodysplasia-related changes are: Complex karyotype (defined as 3 or more chromosomal abnormalities in the absence of 1 of the WHO-designated recurring translocations or inversions, that is, t(8;21), inv(16) or t(16;16), t(9;11), t(v;11)(v;q23.3), t(6;9), inv(3) or t(3;3); AML with BCRABL1); Unbalanced abnormalities: 27 or del(7q); 25 or del(5q); i(17q) or t(17p); 213 or del(13q); del(11q); del(12p) or t(12p); idic(X)(q13); Balanced abnormalities: t(11;16)(q23.3;p13.3); t(3;21)(q26.2;q22.1); t(1;3)(p36.3;q21.2); t(2;11)(p21;q23.3); t(5;12)(q32;p13.2); t(5;7)(q32;q11.2); t(5;17)(q32;p13.2); t(5;10)(q32;q21.2); t(3;5)(q25.3;q35.1).

6. Cases should be classified with the related genetic abnormality given in the diagnosis.

7. The former subgroup of acute erythroid leukemia, erythroid/myeloid type ( $\geq 50\%$  bone marrow erythroid precursors and  $\geq 20\%$  myeloblasts among nonerythroid cells) was removed; myeloblasts are now always counted as percentage of total marrow cells. The remaining subcategory AML, NOS, pure erythroid leukemia requires the presence of  $>80\%$  immature erythroid precursors with  $\geq 30\%$  proerythroblasts.
8. BCR-ABL1 leukemia may present as MPAL; treatment should include a tyrosine kinase inhibitor.
- Table reprinted with permission from the publisher (Döhner *et al.* 2017).

**Table 5. WHO classification of AML with germ line predisposition.** Table adapted with permission from the publisher (Dohner, Estey *et al.* 2017).


<b>Myeloid neoplasms with germ line predisposition without a preexisting disorder or organ dysfunction</b>
<ul style="list-style-type: none"> <li>• AML with germ line CEBPA mutation</li> <li>• Myeloid neoplasms with germ line DDX41 mutation<sup>1</sup></li> </ul>
<b>Myeloid neoplasms with germ line predisposition and preexisting platelet disorders</b>
<ul style="list-style-type: none"> <li>• Myeloid neoplasms with germ line RUNX1 mutation<sup>1</sup></li> <li>• Myeloid neoplasms with germ line ANKRD26 mutation<sup>1</sup></li> <li>• Myeloid neoplasms with germ line ETV6 mutation<sup>1</sup></li> </ul>
<b>Myeloid neoplasms with germ line predisposition and other organ dysfunction</b>
<ul style="list-style-type: none"> <li>• Myeloid neoplasms with germ line GATA2 mutation</li> <li>• Myeloid neoplasms associated with bone marrow failure syndromes</li> <li>• Juvenile myelomonocytic leukemia associated with neurofibromatosis, Noonan syndrome, or Noonan syndrome-like disorders</li> <li>• Myeloid neoplasms associated with Noonan syndrome</li> <li>• Myeloid neoplasms associated with Down syndrome<sup>1</sup></li> </ul>

1. Lymphoid neoplasms also reported.

### 1.2.3 Mutational landscape of AML

Different cancer types carry various numbers of mutations but among cancers, AML is one of the types of cancer with the lowest number of mutations in comparison to other adult cancers (Figure 6) (Alexandrov, Nik-Zainal *et al.* 2013, Lawrence, Stojanov *et al.* 2013). Genetic alterations initially described as karyotypic abnormalities are found in roughly 50% of AML patients by cytogenetic analysis (Dohner, Estey *et al.* 2017, Medinger, Lengerke *et al.* 2016). However, almost half of AML cases do not have any karyotypic abnormalities, and hence are described as cytogenetically normal AML (CN-AML) patients. In CN-AML samples, the mean number of mutations in coding sites, splice sites and RNA genes are around 13 (ranging between 0-51) including eight passenger mutations and five driver mutations (Cancer Genome Atlas Research, Ley *et al.* 2013). So far, numerous mutations have been identified in AML patients, the most frequent of which are depicted in Figure 7. Below, some of the most frequent mutations which cause AML or are important in the development or maintenance of leukemic cells are described.






**Figure 6. The mean number of mutations per genome in various cancer types.** Figure reprinted with permission from the publisher (Grove and Vassiliou 2014).

### ***Nucleophosmin 1 (NPM1) mutations***

The nucleolar protein nucleophosmin 1 plays various cellular functions including protein chaperoning, centrosome duplication, DNA repair, cell proliferation and apoptosis. The most common mutation in AML patients is *NPM1* mutations with a frequency of 25-30% in all AML cases and more specifically 45-64% in CN-AML (Falini, Nicoletti et al. 2007, Schnittger, Schoch et al. 2005). The exact mechanism of *NPM1* in leukemogenesis has not yet been clarified. Since *NPM1* has multiple functions, it is thought that *NPM1* mutations drive leukemia through a combination of cellular processes. Loss of function due to *NPM1* mutations causes uncontrolled centrosome duplication and also promotes myeloproliferation and susceptibility to oncogenic transformation. Moreover, it induces ubiquitination and degradation of TP53 by MDM2. On the other hand, gain of function in cytoplasmic NPM1 inhibits caspase enzymes which blocks apoptosis, induces expression of *Myc* and increases the cell proliferation rate (Heath, Chan et al. 2017).

*NPM1* mutations in AML cases are frequent in the nucleolar localization signal located at the N-terminal and cause abnormal expression of cytoplasmic NPM1 protein rather than in the nucleus. Furthermore, *NPM1* mutations are correlated with *DNMT3A*, *FLT3-ITD* and +8 (40%), *FLT3-TKD* (10-15%) and *IDM* mutations (25%) (Marcucci, Haferlach et al. 2011, Marcucci, Metzeler et al. 2012).



**Figure 7. Common gene mutations in adult AML patient up to 65 years old.** For each gene, the frequency of correlation with other mutations are shown in the boxes. a indicates cohesin genes including RAD21 (~10%), SMC1A (~5%), and SMC3 (~5%). b inv(16)(p13.1;q22) or t(16;16)(p13.1;q22); CBFB-MYH11. c inv(3)(q21.3;q26.2) or t(3;3)(q21.3;q26.2); GATA2,MECOM(EVI1). d TP53 mutations are found in ~45%, and complex karyotypes in ~70% of this class. Figure reprinted with permission from the publisher (Dohner, Estey et al. 2017).

### *Fms-like tyrosine kinase 3 (FLT3) mutations*

FLT3, a tyrosine kinase receptor for the FLT3 ligand was first found with high expression pattern in HSCs and plays an important role in the survival and proliferation of hematopoietic cells (Gilliland and Griffin 2002, Maroc, Rottapel et al. 1993). The most frequent mutations in *FLT3* are in the second tyrosine kinase domain (TKD) or the internal tandem duplication (ITD) in the juxta-membrane domain, which are identified in 20% of all AML patients and 30-45% of CN-AML cases (Kelly, Liu et al. 2002). Gain of function in *FLT3* causes constitutive phosphorylation of some downstream proteins including GAB1, GAB2, SHP-2, AKT, FOXO3a, CBL, STAT5a, ERK1/2, GRB2, SHC, VAV, LYN, GAP, p90RSK, BAD, C/EBPα, and SHI through PI-3 kinase/AKT, RAS/ MAPK, and STAT5 pathways in leukemic cells (Small 2006).

### *Isocitrate dehydrogenase (IDH) mutations*

These enzymes catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. *IDH1* and *IDH2* mutations were identified in 15-20% of all AML patients and 25-30% of CN-AML cases, most often altering the highly conserved arginine residue at codon 132 (R132) of *IDH1*

and at codons R140 and R172 of *IDH2* (Marcucci, Maharry et al. 2010, Patel, Gonen et al. 2012). Moreover, *IDH1* and *IDH2* mutations are also detected more in older AML patients (Paschka, Schlenk et al. 2010).

Mutations of these enzymes in AML, is a gain of function, giving them a new ability to convert  $\alpha$ -ketoglutarate into 2-hydroxyglutarate (2-GH). High levels of 2-GH inhibit TET enzymes, leading to accumulation of histone methylation, subsequently blocking myeloblast differentiation, and increasing stem cell marker expression, suggesting a pro-leukemic effect of *IDH1/2* mutations (Figueroa, Abdel-Wahab et al. 2010, Lu, Ward et al. 2012, Ward, Patel et al. 2010).

### ***DNA methyltransferase 3A (DNMT3A) mutations***

As previously mentioned, the DNA methyltransferase DNMT3A is involved in *de novo* methylation. *DNMT3A* mutations induce global hypomethylation, which is associated with genome instability, and hypomethylation in promoter regions of *HOX* and *MEIS1* genes, which are associated with transcriptional activation (Ferreira, Heyn et al. 2016). *DNMT3A* mutations occur in 18-22% of AML patients and commonly presents as a missense mutation located on R882 compared to other codons in the *DNMT3A* gene (non-R882) (Ley, Ding et al. 2010). *DNMT3A* mutations were detected in the early stages of AML evolution as pre-leukemic mutations and persist during remission (Shlush, Zandi et al. 2014). Yet, a large study of 1700 AML patients revealed that there is no significant correlation between *DNMT3A* mutations and survival (Gaidzik, Schlenk et al. 2013).

### ***Tet methylcytosine dioxygenase 2 (TET2) mutations***

TET2 is a methylcytosine dioxygenase that converts 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC), which plays key role in DNA demethylation. The most common type of *TET2* mutation is loss of function which was found in 9-23% of AML cases (Chou, Chou et al. 2011), however, the prognostic effects of *TET2* mutations remain controversial (Gaidzik, Paschka et al. 2012, Metzeler, Maharry et al. 2011).

*Tet2* mutations in mice are associated with DNA hypermethylation, altered gene expression patterns and subsequent increased self-renewal of HSCs and myeloid expansion (Cimmino, Dolgalev et al. 2017).

### ***Runt-related transcription factor (RUNX1) mutations***

RUNX1, also known as AML1 protein or core-binding factor subunit  $\alpha$ -2 (CBFA2), is a TF with an essential role in normal hematopoiesis (Meyers, Downing et al. 1993). Translocation of *RUNX1* at chromosome 21 with the *ETO* gene on chromosome 8 creates the fusion protein RUNX1-ETO (RUNX1-RUNX1T1) which is frequently found in AML cases (Tang, Hou et al. 2009). The fusion protein RUNX1-ETO suppresses the function of normal RUNX1 in hematopoiesis which induces repression of myeloid-specific genes, causing a block in myeloid cell maturation (Ichikawa, Yoshimi et al. 2013). Moreover, other mutations in *RUNX1* were found in 5-13% of AML cases and were frequently associated with trisomy 13, 21, old CN-AML patients, absence of *NPM1* and poor outcome (Marcucci, Haferlach et al. 2011, Mendler, Maharry et al. 2012, Tang, Hou et al. 2009).

### ***CCAAT enhancer binding protein A (CEBPA) mutations***

CEBPA is a TF which regulates the expression of important genes required for hematopoiesis (Koschmieder, Halmos et al. 2009). *CEBPA* mutations were detected in 6-10% of all AML patients and 15-19% of CN-AML, generally associated with del (9q) (Mrozek, Marcucci et al. 2007). Conditional deletion of *Cebpa* in adult mice blocks CMP to GMP transition and consequently leads to an accumulation of myeloblasts in the BM (Zhang, Iwasaki-Arai et al. 2004). In most AML cases with *CEBPA* mutations, patients often simultaneously carry two mutations (double mutations) commonly known as biallelic in the N-terminal and bZIP regions. Double biallelic mutations, but not single *CEBPA* mutations are associated with favorable overall and higher complete response (CR) (Fasan, Haferlach et al. 2014, Wouters, Lowenberg et al. 2009).

### ***Mixed Lineage Leukemia (MLL1) mutations***


MLL1 (Lysine-specific methyltransferase 2A or KMT2A encodes) is a histone methyltransferase enzyme located on chromosome 11 (11q23). In AML, more than 80 different protein partners were found to generate a fusion oncogene through translocation with *MLL* (Meyer, Hofmann et al. 2013). Leukemia with *MLL* translocations manifest as either acute lymphoid, myeloid or mixed phenotype acute leukemia (MPAL), which are detected in 10% of all leukemia cases (Muntean and Hess 2012). But the incidence of *MLL* rearrangements is 70-80% in infant leukemia patients (Mann, Attarbaschi et al. 2010).

Also, *MLL* rearrangements are common in therapy-related leukemia, typically in patients treated with topoisomerase II inhibitors (e.g., ectoposide, doxorubicin) (Super, McCabe et al. 1993). Translocation of *MLL* is common in AML and it is correlated with poor prognosis. The frequencies of the main partners involved in fusion of *MLL* (*AF4*, *AF9*, *ENL*, *AF10*, *ELL* and *AF6*) in AML are shown in Figure 8. In addition to *MLL* translocations, partial tandem duplication (PTD) of *MLL* is most commonly identified in *de novo* CN-AML and in AML patients with trisomy 11 (Caligiuri, Schichman et al. 1994, Caligiuri, Strout et al. 1998).

*MLL* rearrangements cause upregulation of several *HOX* genes (including *HOXA5*, *HOXA9*, and *HOXA10*) and *MEIS1*, which contribute to the growth advantage and maintenance of self-renewal properties of AML cells (Li, Luo et al. 2009).

### ***Additional sex comb-like 1 (ASXL1) mutations***

ASXL1 is a chromatin-binding protein involved in the epigenetic regulation of cells, yet the clear function of ASXL1 in hematopoiesis is not fully understood. *ASXL1* mutations occur in 5-11% of all AML patients and the mutation rate is five times higher in patients over 60 compared to those under 60 years old (Metzeler, Becker et al. 2011). *ASXL1* mutations cause loss of H3K27me3 mediated by the PRC2 complex and upregulated expression of *HOXA* genes. *Asxl1* mutations in combination with NRasG12D expression promotes myeloid leukemogenesis *in vivo* (Abdel-Wahab, Adli et al. 2012).



**Figure 8. The Frequency of MLL-rearranged AML with main seven fusion partners.** The right pie charts show the breakdown of the relative frequencies of MLL fusion partners based on age group. Figure reprinted with permission from the publisher (Winters and Bernt 2017).

### ***Tumor protein p53 (TP53) mutations***

TP53 regulates quiescence and self-renewal of HSCs, and so *p53* mutations promote HSPC proliferation, leading to an accumulation of DNA damage and ultimately transformation of HSCs into pre-leukemic cells (Prokocimer, Molchadsky et al. 2017). Mutations of *TP53* are reported in 8-14% of AML cases and *TP53* alterations are most often associated with a complex karyotype, poor prognosis and resistance to chemotherapy (Haferlach, Dicker et al. 2008, Rucker, Schlenk et al. 2012).

### ***KIT mutations***

KIT tyrosine kinase is a transmembrane receptor for mast cell growth factor (MGF) and critical for normal hematopoiesis (Sattler and Salgia 2004). The *KIT* mutation frequency is less than 5% in all AML patients, but identified in 22-29% of AML cases with CBF mutations (Paschka, Marcucci et al. 2006). Moreover, overexpression of *KIT* is identified in 60-80% of AML patients (Kindler, Breitenbuecher et al. 2004). *KIT* mutations are associated with a high relapse rate and shorter overall survival (Boissel, Leroy et al. 2006).

*KIT* mutations trigger constitutive KIT kinase activation, promoting downstream signaling pathways which contribute to factor-independent growth of hematopoietic cells and induce leukemogenesis (Lennartsson, Jelacic et al. 2005).

### **1.2.4 Prognosis factors**

A crucial component of AML treatment is an accurate assessment of AML prognosis. Prognostic factors can provide information for physicians to decide on a treatment strategy between standard or increased intensity treatment, consolidation chemotherapy or transplantation. Prognostic factors can be divided in two categories: patient-related, such as age and performance status and disease-associated factors including white-cell count, cytotoxic therapy for another disorder or prior myelodysplastic syndrome, cytogenetic and molecular aberrations (Bullinger, Dohner et al. 2017, Medinger, Lengerke et al. 2016, Meyer and Levine 2014).

In 2010, the European LeukemiaNet (ELN) proposed a standardized classification of risk stratifications in adult AML patients according to the correlation of cytogenetic and molecular genetics with clinical data (Dohner, Estey et al. 2010). In the updated 2017 classification, patients are divided into three risk groups: favorable, intermediate and adverse.

(Table 6) (Dohner, Estey et al. 2017). Recently, the new techniques such as FISH, polymerase chain reaction (PCR) and gene sequencing have provided tools to enable a more comprehensive view of AML prognosis.

In abnormal karyotype patients, alterations of chromosome 5, 7, 11q23 and more than three chromosomal abnormalities are associated with poor prognosis and reduced survival, whilst t(8;21)(q22;q22), t(15;17)(q22;q12) and inv(16)(p13.1;q22) abnormalities are associated with longer survival (Dohner, Weisdorf et al. 2015, Medinger and Passweg 2017).

*KIT* mutations in patients with t(8;21) translocation increase the risk of relapse and decreased overall survival (Estey 2014, Qin, Zhu et al. 2014). The presence of *TP53* or *DNMT3A* mutations are associated with poor prognosis (Kihara, Nagata et al. 2014, Shivarov, Gueorguieva et al. 2013). In CN-AML, partial tandem duplication of *MLL* is associated with a worse prognosis (Patel, Gonen et al. 2012).

Several studies have shown that *FLT3-ITD* mutations in older CN-AML patients (>60 years old) are associated with poor outcome (Patel, Gonen et al. 2012, Port, Bottcher et al. 2014). ITD of *FLT3* is associated with a poor prognosis and high relapse rate (Kelly, Liu et al. 2002, Meyer and Levine 2014, Saultz and Garzon 2016), constitutively activates tyrosine kinase and subsequently activates RAS, MAPK and STAT5 pathways and induces cell proliferation (Gale, Green et al. 2008, Kayser, Schlenk et al. 2009, Kelly, Liu et al. 2002).

**Table 6. Risk assessment of AML according European LeukaemiaNET (ELN) 2017.** Table adapted with permission from the publisher (Medinger and Passweg 2017).

Risk category	Genetic abnormality
Favorable	<ul style="list-style-type: none"> <li>t(8;21)(q22;q22); RUNX1-RUNX1T1</li> <li>Inv(16)(p13;q22) or t(16;16)(p13;q22); CBFB-MYH11</li> <li>Mutated NPM1 without FLT3-ITD or with FLT3-ITD<sup>low</sup> (allelic ratio &lt;0.5)</li> <li>Biallelic mutated CEBPA</li> </ul>
Intermediate	<ul style="list-style-type: none"> <li>Mutated NPM1 and FLT3-ITD<sup>high</sup> (allelic ratio ≥0.5)</li> <li>Wild-type NPM1 without FLT3-ITD or with FLT3-ITD<sup>low</sup> (without adverse-risk genetic lesions)</li> <li>t(9;11)(p21;q23); MLLT3-KMT2A</li> <li>Cytogenetic abnormalities not classified as favorable or adverse</li> </ul>
Adverse	<ul style="list-style-type: none"> <li>t(6;9)(p23;q34); DEK-NUP214</li> <li>t(v;11q23); KMT2A rearranged</li> <li>t(9;22)(q34;q11); BCR-ABL1</li> <li>inv(3)(q21;q26) or t(3;3)(q21;q26); GATA2, MECOM (EV11)</li> <li>-5 or del (5q); -7; -17/abn(17p)</li> <li>Complex karyotype (three or more unrelated chromosome abnormalities, monosomal karyotype (defined by the presence of 1 single monosomy))</li> <li>Wild-type NPM1 and FLT3-ITD<sup>high</sup></li> <li>Mutated RUNX1</li> <li>Mutated ASXL1</li> <li>Mutated TP53</li> </ul>

*NPM1* mutations are associated with sensitivity to intensive chemotherapy, which may result in improved outcomes in both young and older patients (Dohner, Estey et al. 2017, Dohner, Schlenk et al. 2005).

As a result, these three markers *NPM1* mutation, *CEBPA* mutation, and *FLT3* ITD are currently used in clinical practice (Dohner, Estey et al. 2010). Therefore, deep genetic analyses of AML patient samples are critical for prognostic classification and treatment plan strategies, however the ability to predict resistance to treatment and relapse remains limited (Walter, Othus et al. 2015).

### **1.2.5 The bone marrow niche of AML**

Similar to normal HSCs, leukemic cells also need a proper niche in which to reside and expand. Recent studies have confirmed that the BM niche supports development and expansion of hematopoietic malignancies. Massive alterations of the BM microenvironment with age, in addition to mutation accumulation, contribute to leukemogenesis. For example, activation of the osteoblastic cell-specific parathyroid hormone (PTH) receptor enhanced frequency of LSCs in the *MLL-AF9* AML mouse model (Krause, Fulzele et al. 2013).

Interestingly, there is also evidence that alterations to the BM niche can induce AML. For instance, deletion of *Dicer1* or *Sbds* in osteoprogenitors induced myelodysplasia and predisposition to AML (Raaijmakers, Mukherjee et al. 2010). Additionally, activating mutation of  $\beta$ -catenin in osteoblast cells induces high activation of Notch signaling in progenitor cells, resulting in an altered differentiation of myeloid and lymphoid cells, and ultimately inducing AML (Kode, Manavalan et al. 2014). Altogether, these studies indicate that alterations to the BM niche can provide appropriate conditions which promote hematopoietic malignancies.

Further studies have demonstrated that the BM niche is altered during development of CML-blast crisis resembling AML by induction signals from the leukemic cells. It has been shown that leukemic cells in direct contact with MSCs secrete TPO and CCL3 that stimulate MSCs to overproduce functionally altered osteolineage cells which support LICs (Schepers, Pietras et al. 2013).

In summary, it seems that the BM niche and leukemic cells have mutual cross-talk during initiation, development and AML relapse.



### 1.2.6 Clonal hematopoiesis and AML evolution

Since HSCs divide during the whole human life, they may acquire random mutations with time which accumulate and are passed on to the next generation of cells. Although most mutations in HSCs are benign, mutations that induce enhanced self-renewal capacity or proliferation of HSCs and/or reduce apoptosis, cause clonal advantage.

The two-hit model was first proposed by Gilliland and Griffin (Gilliland and Griffin 2002) for clonal hematopoiesis (CH) and subsequently leukemogenesis. In this model two classes of mutations are required. Class I mutations contribute to the activation of proliferation pathways or to survival advantage such as *FLT3-ITD*, *K/N-RAS*, *TP53* and *c.KIT*. Whereas class II mutations impair normal differentiation of hematopoietic cells such as *NPM1*, *CEBPA*, *PML/RAR $\alpha$*  and *MLL* fusion genes. In addition to these two classes of mutations, there are some mutations of epigenetic factors such as *DNMT3A*, *TET2*, *IDH1* and *IDH2* which effect both proliferation and differentiation, and so can be classified in to a third group (De Kouchkovsky and Abdul-Hay 2016).

Additionally, other mechanisms such as the immune system, BM niche alterations and selective pressure on somatic variants (aging, chemotherapy and exogenous stress such as smoking) also influence CH (Bowman, Busque et al. 2018). The most common mutations present in CH are in epigenetic factors like *TET2*, *DNMT3A* and *ASXL1* and splicing factors such as *SF3B1*, *SRSF2* and *U2AF1* (Bowman, Busque et al. 2018). These mutations in CH prime them to develop hematopoietic malignancies.

Mutations that contribute to cancer initiation and growth are called driver mutations, whilst those that do not affect cancer growth are termed passenger mutation. However, the classification of cancer depends on its microenvironment. For example, certain AML mutations can confer drug resistance so that during cancer treatment sub-clones carrying these mutations can survive and expand, whereas in the absence of treatment these mutations might have a growth disadvantage effect (Grove and Vassiliou 2014). On the other hand, some passenger mutations may be selected in a specific microenvironment condition and induce drug resistance or growth advantage.

Cancer evolution similar to Darwinian natural selection was first suggested by Peter Nowell (Nowell 1976). Emerging data from AML patient samples using new technologies, particularly NGS, are providing a better view of mutational frequencies, clonal evolution of the disease, sub-clonal patterns and into the epigenetic landscape of AML. The result from a clonality analysis of 50 whole genome sequences and 150 whole-exomes sequences of AML patient samples revealed that the majority of AML cases are clonally heterogeneous with a

founding clone and at least one sub-clone at the diagnosis stage (Cancer Genome Atlas Research, Ley et al. 2013). Importantly, two major clonal evolution patterns during relapse were determined: either a founding clone or a surviving sub-clone acquired additional mutations most likely contributed to drug resistance and expanded at relapse stage (Ding, Ley et al. 2012).

New findings from NGS of pre-leukemic stem cells and clonal evolution of AML revealed that pre-leukemic HSCs already contain recurrent common mutations in epigenetic regulators, such as *DNMT3A*, *ASXL1*, *IDH1*, *IDH2*, and *TET2*. They are therefore thought to occur in the early stage of AML evolution (Corces-Zimmerman, Hong et al. 2014, Kronke, Bullinger et al. 2013, Shlush, Zandi et al. 2014). Moreover, the frequency of somatic mutations (*DNMT3A*, *TET2*, and *ASXL1*) were rare in individuals under 40 years old but increased with age and the presence of these mutations was associated with a higher risk of hematologic cancers (Busque, Patel et al. 2012, Genovese, Kahler et al. 2014, Jaiswal, Fontanillas et al. 2014). Other somatic mutations which are known in AML are *FTL3*, *NPM1*, *WT1*, *RUNX1*, *RTRP*, *PHF6* and *ETV6* (Zhang, Lv et al. 2016).

### **1.2.7 Leukemic stem cells**

A small population of cells amongst AML blast cells, have limitless self-renewal capacity and the ability to propagate and go on to develop AML. These cells are known as leukemic stem cells (LSCs) or leukemia initiating cells (LICs). LICs are a rare population (roughly 1 per  $10^6$  bulk) in many AML patients capable of initiating leukemia in immunodeficient mice and maintaining long-term in serial transplanted mice with self-renewing capacity. There is some evidence showing that LICs are not only originated from HSCs, but certain mutations in differentiated cells may also transform them in to LSCs (Cozzio, Passegue et al. 2003, Huntly, Shigematsu et al. 2004).

The critical step of LIC detection is to define markers which are associated with functional properties of LICs. LICs usually exist in the  $CD34^+CD38^-$  fraction of AML cells in patients (Lapidot, Sirard et al. 1994). However in more than 50% of samples LICs reside in  $CD34^+CD38^+$  or even the  $CD34^-$  population (Kreso and Dick 2014). Further experiments revealed in 80% of AML patients LSCs exist in both fractions; LMPP-like (lymphoid-primed multipotent progenitors) LSCs ( $Lin^-CD34^+CD38^-CD90^-CD45RA^+$ ) and GMP-like LSCs ( $Lin^-CD34^+CD38^+CD123^+CD45RA^+$ ). Interestingly, LMPP-like LSCs could give rise to GMP-like LSCs but not the converse (Goardon, Marchi et al. 2011).

From a clinical perspective, to achieve long-term complete remission (CR), it is critical to eliminate LICs. However, there are several obstacles to target LICs with conventional therapies. First, there is no standard method to characterize LICs which demonstrate their properties. Secondly, LICs are a rare population and quiescent so it is difficult to eliminate them with current drugs which target proliferative bulk cells. Thirdly, LICs are usually resistant to apoptosis and they highly express channels for drug efflux (Thomas and Majeti 2017). In addition, recent studies have shown heterogeneity in LICs both between and within individual patients (Eppert, Takenaka et al. 2011, Goardon, Marchi et al. 2011). Thus, it is needed to identify new targets against LICs to develop more effective therapies.

### **1.2.8 AML treatment**

#### **Current therapy**

The common therapeutic approach in AML patients not much has changed since 1970 (Dohner, Estey et al. 2010). The first line of treatment is to assess whether intensive chemotherapy is suitable for a AML patient. After intensive therapy, post-induction therapy is important to achieve CR. Two types of post-induction strategies are: conventional chemotherapy and hematopoietic cell transplantation.

#### **Induction therapy**

The aim of induction therapy is to diminish AML cells and achieve CR. The standard induction therapy is chemotherapy. Intensive induction therapy is applicable to young adult AML patients under 60 years old and older patients specifically with *NPM1* mutations and CBF leukemia (Saultz and Garzon 2016). The standard chemotherapy protocol is the “7+3” regimen with a combination of continuous infusion of cytarabine (100 mg/m<sup>2</sup> daily for one week) and an anthracycline (either daunorubicin 60-90 mg/m<sup>2</sup> or idarubicin 10-12 mg/m<sup>2</sup> on days 1, 2 and 3). The current conventional therapies for AML patients are summarized in Table 7.

The CR rate is 65-73% in adult patients younger than 60 years and 38-62% in patients over 60 years old (Estey and Dohner 2006, Fernandez, Sun et al. 2009, Lowenberg, Ossenkoppele et al. 2009). CR is defined as (1) <5% blasts in a BM aspirate sample with marrow spicules (no blasts with Auer rods or persistence of extramedullary disease), (2) absolute neutrophil count (ANC) >1000/μl, and (3) platelets ≥100,000/μl (Cheson, Bennett et al. 2003).

## Consolidation (post-induction) therapy

The aim of consolidation strategies is to eradicate minimal residual disease (MRD), prevent relapse and ideally to achieve a complete cure with either chemotherapy or transplantation. MRD can be monitored by using molecular based approaches such as real-time quantitative PCR (RT-qPCR) or target gene sequencing to detect genetic targets specific to AML cells and/or multi-parameter flow cytometry (MFC) to read-out leukemic cells based on their immune phenotype (Grimwade and Freeman 2014).

In consolidation chemotherapy, intermediate dose of cytarabine ( $1-1.5\text{g/m}^2$ ) is used twice daily on days 1, 3 and 5 in three or four cycles in adults younger than 60 years old, where the cure rate is 60-70% (Byrd, Mrozek et al. 2002). However, in patients older than 60 years, the cure rate is dismal, between 10-15%, therefore new strategies of maintenance therapies need to be investigated (Dohner, Weisdorf et al. 2015).

Hematopoietic stem cell transplantation is a treatment strategy for patients who are unlikely respond to the conventional induction therapy and to extend CR. Prior to transplantation, chemoradiotherapy is chosen to deplete leukemic cells, suppress the immune system, and to

**Table 7. Summary of current conventional care of AML patients.** Table adapted with permission from the publisher (Dohner, Weisdorf et al. 2015).

Type of therapy	Age	Regimen
Induction therapy	Patients 16–60 yr	3 days of an intravenous anthracycline (daunorubicin $60\text{ mg/m}^2$ ; idarubicin $10-12\text{ mg/m}^2$ ; mitoxantrone $10-12\text{ mg/m}^2$ ) and 7 days of continuous-infusion cytarabine ( $100-200\text{ mg/m}^2$ ) (“3+7” induction)
	Patients >60 yr	For patients with favorable-risk and intermediate-risk cytogenetic findings and no coexisting conditions, induction therapy is the same as that in younger patients, and dose reduction may be considered for individual patients
Consolidation therapy	Patients 16–60 yr	Patients with favorable genetic risk (according to ELN) should Receive 2-4 cycles of intermediate-dose cytarabine ( $1-1.5\text{ g/m}^2$ intravenously, usually administered every 12 hr over 3 days, or $1-1.5\text{ g/m}^2$ intravenously on days 1-6); for patients with intermediate-I, intermediate-II, or adverse risk, allogeneic hematopoietic-cell transplantation should be strongly considered; if not possible, consolidation therapy should be administered as above; combination chemotherapy (e.g., mitoxantrone-cytarabine) may be superior in patients with adverse-risk AML
	Patients >60 yr	Patients with favorable ELN genetic risk (less common) and no coexisting conditions should receive 2-3 cycles of intermediatedose cytarabine ( $0.5-1\text{ g/m}^2$ intravenously, every 12 hr on days 1-3, or $0.5-1\text{ g/m}^2$ intravenously, on days 1-6)

provide space for donor cells in the BM. The most common regimen for chemoradiotherapy is fludarabine in combination with cyclophosphamide or other alkylating reagents (such as melphalan and busulfan) and total-body irradiation. The key dilemma after transplantation is graft-versus-host-disease (GVHD), in which donor immune cells recognize the host cells as foreign and attack the recipient's organs. Therefore, accurate HLA matching is required to decrease the risk of GVHD.

Despite new techniques such as NGS to predict prognosis and to improve treatment strategy selection, the heterogeneity of AML and the varied genetic landscape of sub-clones make it difficult to estimate the risk of relapse.

### **Novel treatment strategies**

Since AML is a heterogeneous disease within patients and even within sub-clones, the standard chemotherapy strategy is not the best option for all patients and given that chemotherapy does not specifically target leukemic cells, it is known to cause severe side effects. Personalized medicine (also known as precision medicine) is the new goal in the cancer field, which refers to individual patient based-strategies, where their genetic background, prognosis and predicted response and treatment strategies are tailored to each individual patient. However, it is demanding to design new drugs or inhibitors specifically targeting driver genes in cancer cells for treatment based on the mutational background to improve cure rate, increase survival rate and prevent relapse. Here, new inhibitors and treatment strategies in AML are briefly described.

### **FLT3 Inhibitors**

The first generation of multiple kinase inhibitors such as midostaurin, lestaurtinib, tandutinib, sunitinib and sorafenib, have recently shown transient reduction of leukemic cells and increased toxicity (Sudhindra and Smith 2014). The second generation of *FLT3* inhibitors including quizartinib and crenolanib are being used in the clinical phase and have better potency with less side effects. However, drug resistance is a main obstacle when using single inhibitor for FLT3 mutations. Recently, the drug Midostaurin was approved by the FDA to be used in combination with daunorubicin and cytarabine for *FLT3* mutation positive AML patients. Additionally, G-749 and ASP2215, as novel *FLT3* inhibitors with a decreased risk of drug resistance, have recently been used in clinical trials (Lee, Kim et al. 2014).

## **IDH Inhibitors**

In 20% of AML samples *IDH-1* or *IDH-2* have gain of function mutations (Cancer Genome Atlas Research, Ley et al. 2013). There are some *IDH1* and *IDH2* inhibitors currently in phase I and II clinical trials, such as NCT02381886, NCT0195498 and NCT02074839, AG-221 and AG-120, CB-839, Erwinaze (Fathi, Wander et al. 2015, Medinger and Passweg 2017). The response rates in relapsed patients to AG-221 and AG-120 was 40% and 31% respectively.

## **HDAC inhibitors**

Epigenetic modifiers HDACs catalyze deacetylation of histones and involve in gene silencing. Mutations or dysregulations of many acetyltransferases and deacetylases have been identified in leukemia including p300, CBP, AML1. Therefore, HDAC inhibitors including vorinostat, mocetinostat and SAHA appeared as an attractive therapeutic strategy for AML. Since the response rate in monotherapy studies using single HDAC inhibitors were low (13-17%), the current trial strategies focus on combination of HDAC inhibitors with chemotherapy or other epigenetic inhibitors (Saygin and Carraway 2017).

## **STAT inhibitors**

Deregulation of STAT signaling in AML is associated with an increased expansion, apoptosis block and abnormal differentiation of leukemic cells. Constitutive activation of STAT3 and STAT5 either alone or together has been reported in AML patients (Bar-Natan, Nelson et al. 2012). Furthermore, *STAT3* tyrosine phosphorylation is associated with a worse prognosis and was found to be upregulated in 50% of AML patients (De Kouchkovsky and Abdul-Hay 2016). Moreover, overexpression of STAT3 signaling contributes to *FLT3* inhibitor resistance in AML cells (Zhou, Bi et al. 2009). Therefore, *STAT3* inhibitors such as OPB-31121 could be used for patients who are being treated with *FLT3* inhibitors.

## **MLL inhibitors**

There are many approaches to target MLL fusion proteins in AML patients with MLL translocations. One target is Disruptor of telomeric silencing 1-like (DOT1L), a histone methyltransferase, which is recruited by most of the MLL fusion proteins. Abnormal interaction of MLL oncogene and DOT1L results in methylation of H3K79 and activation of

MLL downstream target genes including *HOXA9* and *MEIS1* (Kavanagh, Murphy et al. 2017). Other inhibitors against Menin, Polycomb proteins, LSD1 and Bromodomain proteins can be used to target MLL-rearranged leukemias (Winters and Bernt 2017).

### **Nuclear Exporter Inhibitors**

Chromosome region maintenance 1 (CRM1) is a nuclear exporter protein which is responsible for exporting several tumor suppressor proteins such as P53, P21, P73, FOXO1, RB1 and NMP1 (Fukuda, Asano et al. 1997). Overexpression of CRM1 causes exorbitant transportation of tumor suppressor proteins from the nucleus to the cytoplasm, which is associated with poor prognosis and decreased survival (Kojima, Kornblau et al. 2013). Selinexor as a CRM1 inhibitor is in clinical trials to be used as a drug for AML patients with high CRM1 expression levels (Etchin, Sun et al. 2013, Ranganathan, Yu et al. 2012).

### **Clofarabine**

Clofarabine is a second-generation purine nucleoside analogue FDA-approved for treating relapsed or refractory childhood ALL patients. Clofarabine enters cells via passive or active transport and is transformed to its active triphosphate form by kinases. Clofarabine inhibits both DNA polymerase and ribonucleotide reductase, which in turn inhibits DNA replication as well as RNA transcription (Tiley and Claxton 2013). Treatment of older AML patients with Clofarabine resulted in a 40% overall response rate (Burnett, Russell et al. 2013, Kantarjian, Erba et al. 2010). Combination of Clofarabine with a low dose of cytarabine significantly increased the CR rate from 31% to 63% in patients over the age of 60 (Faderl, Ravandi et al. 2008).

### **Immune and Cell Therapies**

A new strategy of induction therapy is immune therapy using an anti-CD33 antibody, which in testing showed a similar response rate compared to chemotherapy treatment but reduced risk of relapse and increased survival. The transmembrane receptor CD33 is expressed on cells of myeloid origin but not on normal HSCs. Widespread expression of CD33 in AML cells promoted the idea of targeting CD33 to treat AML patients (Appelbaum and Bernstein 2017). A humanized anti-CD33 monoclonal antibody (Gemtuzumab ozogamicin) conjugated with the cytotoxic agent calicheamicin was tested in five randomized trials and approved by

the FDA in 2000 (Hills, Castaigne et al. 2014). Although further clinical trials in 637 patients failed to show improvements in CR or overall survival (Petersdorf, Kopecky et al. 2013), recent trials showed a benefit in survival and relapse rates in old AML patients not eligible for intensive chemotherapy and also in patients with an intermediate to favorable cytogenetic-risk pattern (Amadori, Suci et al. 2016).

Another new cell therapy strategy is to use engineered T cells called chimeric antigen receptor (CAR)-transduced T cells (CARTs) to express antigen receptors against specific cell surface antigens in target AML cells. One type of CARTs in preclinical trials is against CD123, which is expressed on the majority of AML cells (Gill, Tasian et al. 2014). CD33-specific CART cells showed improved survival in a xenograft AML mouse models (Kenderian, Ruella et al. 2015). New therapeutic agents in ongoing clinical trials for AML treatment are listed in Table 8.

**Table 8. Summary of new therapeutic agents in ongoing clinical trials.** Table adapted with permission from the publisher (Yu and Zheng 2017).


Agent	Name	Target	Phase of testing
Molecular antibody	Gemtuzumab ozogamicin	CD33	I, II, III
	Vadastuximab talirine (SGN-CD33A)	CD33	I, II
	AMG 330	CD33	I
	HuM195	CD33	I, II
	Yttrium Y 90 anti-CD45 monoclonal antibody BC8 (90Y-BC8)	CD45	NA
	KB004	EphA3	I, II
	Ipilimumab	CTLA-4	I
	Brentuximab	CD30	I, II
	Ulocuplumab	CXCR4	I
FLT3 inhibitors	Lestaurtinib	FLT3	I, II
	Midostaurin	FLT3	I, II, III
	Sorafenib	FLT3	I, II, IV
	Quizartinib	FLT3	I, II, III
	Crenolanib	FLT3	I, II, III
	Gilteritinib	FLT3	II, III
	Pexidartinib (PLX3397)	FLT3	I, II
AURK inhibitors	Alisertib	AURKA	I, II
mTOR kinase inhibitors	Sirolimus	mTOR	I, II
	Temsirolimus	mTOR	I, II
	Everolimus	mTOR	I, II
Epigenetic agents	Decitabine	Methyltransferase	I, II, III
	Azacitidine	Methyltransferase	I, II
	Vorinostat	Histone acetylase	I, II
	Panobinostat	Histone acetylase	I, II
CAR-T cell therapy	CAR-T33	CD33	I, II
	CAR-T123	CD123	I, II



### 1.2.9 Childhood AML

Among all AML cases, 6% of them develop in childhood (Dores, Devesa et al. 2012) and 20% of pediatric leukemic cases are AML (Yu and Zheng 2017). The survival rate of AML in patients under 20 years old have improved during the last 40 years from 20% in 1980 to 70% in 2012 of AML cases, but the clinical outcomes of childhood AML in many subtypes remains low (Gamis, Alonzo et al. 2013, Rubnitz and Inaba 2012, Rubnitz, Inaba et al. 2014).

Certain mutations common in adult AML are rare in childhood AML and vice versa, suggesting differing mechanisms and pathogenesis of AML in adults and children. The main molecular aberrations in childhood AML are depicted in Figure 9. The major differences between childhood AML compared to adult AML are a significantly higher frequency of *MLL*-related leukemia, particularly in patients under two years old, and a lower incidence of cytogenetic abnormalities (Cooper, Franklin et al. 2012). Therefore, the role of molecular risk factors such as *FLT3-ITD*, *IDH1* and *DNMT3* mutations is more prominent in childhood AML cases (Ho, Alonzo et al. 2010, Ho, Kutny et al. 2011, Meshinchi, Alonzo et al. 2006). Furthermore, pediatric AML patients are typically able to tolerate more intensive therapy, which affects the overall outcome compared to adult AML patients.



**Figure 9. Frequency the main cytogenetic (A) and molecular (B) abnormalities identified in childhood AML younger and older that 2 years old.** Figure reprinted with permission from the publisher (Masetti, Vendemini et al. 2015).



## **2 AIM OF THE THESIS**

The overall aim of the thesis was to identify and characterize novel effectors and pathways important in maintenance of AML and in normal hematopoiesis.

The specific aims were:

### **Study I:**

Identify and characterize novel target genes that are specifically required for the maintenance of AML but not normal hematopoietic cells.

### **Study II:**

Identify and characterize novel transcriptional regulators important in the maintenance and disease progression of AML.

### **Study III:**

To investigate the role of the epigenetic enzyme *EHMT1* in AML.

### **Study IV:**

To investigate if targeting SAMHD1 represents an attractive therapeutic approach to potentiate the effects of ara-C treatment in hematological malignancies.

### **Study V:**

To determine the role of the epigenetic factor *NAPIL3* in HSCs and hematopoiesis.



### 3 METHODOLOGICAL APPROACHES

#### 3.1 LENTIVIRAL TRANSFECTION AND TRANSDUCTION

For lentivirus production, The adherent 293FT cells were co-transfected with the vectors, pAX8 and pCMV-VSVG, using the calcium phosphate transfection method (Kingston, Chen et al. 2003). Briefly, the 293FT cells were split in 10 cm dishes on the day before transfection and the cells fed with 9 ml DMEM media containing 10% FBS 4 hours before transfection. For the transfection 10 µg vector DNA, 8 µg pAX8 and 4 µg pCMV-VSVG were mixed with 64 µl of 2 M CaCl<sub>2</sub> and water was added up to 500 µl. 500 µl HEPES was added into a 5 ml FACS tube and the DNA/CaCl<sub>2</sub> solution added dropwise with a pipette whilst simultaneously aerating the mixture by constantly bubbling with a mechanical pipettor attached to a 1 ml pipette and then the mixed solution was immediately vortexed for 5 seconds. The precipitate was incubated for 10 minutes at room temperature and 1 ml of the mixed solution was added to a 10 cm plate, covering almost everywhere and incubate for 6 hours or overnight. The medium was replaced with 6 ml of medium suitable for the target cells for transduction, and the transfected plates were incubated at 32°C to produce virus. Virus supernatant was harvested 24 h and 48 h post transfection and was concentrated by centrifugation at 6000 x g for 16 h at 4°C.

The AML cells and normal hematopoietic cells were transduced with the virus supernatant by spinoculation, i.e. by centrifugation of a mixture of cells and virus for 2 h at 1000 x g in 6 or 12 well non-tissue treated plates. Transduced cells were selected by adding puromycin at the final concentration of 2 µg/ml for 48 h.

#### 3.2 LARGE SCALE SHRNA SCREEN

RNA interference (RNAi) was first discovered in *Caenorhabditis elegans* as double stranded RNA-mediated gene silencing. Double stranded RNA (dsRNA) is a substrate for Dicer (RNase III enzyme) that produces 20-25 bp dsRNA, which is recognized by the RNA-induced silencing complex (RISC). RISC destroys the “passenger strand” and keeps the “guiding strand”. The RISC complex and guiding strand bind to target mRNA and digest it, resulting in post-transcriptional gene silencing (Gao, Yang et al. 2014).

The major challenge in molecular biology is to find out the functional role of genes in specific strains of cells. RNAi technology (Fire, Xu et al. 1998) is a tool to silence target genes and investigate the role of the gene. The advantage of RNAi is the ability to investigate the role

of a gene based on the phenotype-associated loss of function. Therefore, RNAi has become the appropriate method for functional genomics, signal transduction, and drug target discovery assays. In the cancer field, using RNAi technology helps to identify oncogenes which are essential for cancer cell survival and to find target genes for cancer treatment. Also, it is possible to pool individual RNAi and make libraries for genome-wide or high-throughput screens (HTS) to quickly and simultaneously screen multiple genes. Firstly, the readout of RNAi HTS should be well-defined such as proliferation, apoptosis or cell morphology. Similarly, a proper and easy phenotypic assay should be defined based on the readout.

We used two libraries: library module 1 including 27,500 shRNAs (short hairpin RNAs) against 5,000 signaling pathway genes for study II and library module 2 including 27,500 shRNA targeting disease-associated genes in study I. Each individual shRNA contained a unique barcode.


In studies I and II, we used lentiviral vector-based shRNA screens. To ensure we have enough cells per individual shRNA ( $>1000$  cells/shRNA), we cultured approximately  $3 \times 10^8$  of several AML cell lines (THP-1, NOMO-1, and the mouse MLL-AF9 AML) with a concentration of 250,000 cells/ml which were transduced with pooled lentiviral shRNA libraries. To minimize the number of cells with multiple viral integrations, we kept the transduction efficiency at less than 25%, as determined by flow cytometry to measure RFP<sup>+</sup> cells. After 48 h of transduction, we added puromycin (2 $\mu$ g/ml) to select transduced cells, after which, half of these were collected for an initial time point (T0) where the selection control cells were completely dead. The rest of the cells were harvested after 10 cell divisions (T10) in culture. Genomic DNA was extracted from T0 and T10 cells by QIAgene DNA extraction kit according to the manufacturer's instructions, and the barcodes were amplified from the extracted genomic DNA using nested PCR according to Collecta's protocols. PCR-amplified barcodes were purified using the QIAgene PCR purification kit and used for NGS (HiSeq 2000, Illumina). Barcodes were deciphered using Collecta's software to detect the number of reads per barcode, which represented the frequency of individual shRNAs targeting specific genes at T0 and T10. After normalization of reads between samples, the ratios of individual barcodes were calculated by dividing T10/T0.

There are three possibilities for each shRNA according the ratio number. If the ratio of T10/T0 is  $>1$ , it means that the target gene was a growth suppressor and by knocking down the gene the cells containing the shRNA proliferate more. The second scenario is that T10/T0 is equal to 1, meaning that the target gene did not influence cell growth and finally, the third possibility is that ratio of T10/T0  $<1$ , meaning that the target gene has an important role in

cell growth. We selected the shRNA targeting genes from the third scenario with at least 5-fold reduction ( $T_{10}/T_0 < 0.2$ ) in AML cell growth for further validations.

### 3.3 CRISPR/CAS 9 GENOME EDITING


The clustered regularly interspaced short palindrome repeats (CRISPR)/Cas9 technology is a new gene editing system. CRISPR/Cas9 acts as an adaptive immune system in bacteria for cleaving foreign genetic elements such as viruses and plasmids via RNA-guided nucleases (Deveau, Garneau et al. 2010). Cas9 (a non-specific CRISPR-associated endonuclease 9) is guided by small RNA “gRNAs” to the target DNA sequences. CRISPR systems have been divided into two major classes based on their components and mechanisms of action. Class 1 including type I, III and IV involve several effector proteins but class 2 (type II, V and VI) only requires one RNA-guided endonuclease (Wang, La Russa et al. 2016). Cas9 is guided to target DNA by two hybridized RNAs: crRNAs, that identifies target DNA through a 20 base pair (bp) recognition sequence and tracrRNAs, which hybridizes with the crRNAs. To simplify the system for genome editing, crRNA and tracrRNA, can be engineered as a single chimeric guide RNA (sgRNA) (Jinek, Chylinski et al. 2012). The most widely used Cas9-based genome editing system is a single protein Cas9 with a single RNA sgRNA (Cas9-sgRNA) which binds to 20 bp target DNA adjacent to a protospacer adjacent motif (PAM) sequence (Figure 10).



**Figure 10. Schematic of the RNA-guided Cas9 nuclease.** protospacer adjacent motif (PAM), non-homologous end joining (NHEJ), homology-directed repair (HDR). Figure reprinted with permission from the publisher (Cui, Xu et al. 2018).

CRISPR/Cas9 tools edit the genome via two mechanisms: non-homologous end joining (NHEJ) or homology-directed repair (HDR) (Figure 9). In the absence of a repair template, the Cas9 enzyme creates DNA double strand breaks (DSBs) and re-ligates through the error-prone NHEJ process, often producing an insertion/deletion or single nucleotide mutation, which result in frameshifts and eventually gene knockout. HDR is another DNA repair pathway which uses a repair template which can be used to induce precise and defined gene modification at a target site.

Lastly, since CRISPR/Cas9 is flexible and robust technique, it is becoming attractive tool to use not only for genome editing but for various genome modification applications including gene regulation (silencing or activation), epigenome editing, chromatin imaging, chromatin topology, RNA targeting (Figure 11).



**Figure 11. Main applications of CRISPR/Cas9 technology.** Figure reprinted with permission from the publisher (Adli 2018).

For CRISPR/Cas9-mediated knockout studies, gRNAs were designed from Optimized CRISPR Design - MIT (<http://crispr.mit.edu>) provided by the Zhang laboratory and were cloned into the vector for inducible expression pRSITEP-U6Tet-(sh)-EF1-TetRep-2A-Puro expressing fluorescent marker (GFP, BFP, or iRFP670).

First, we developed a stable Cas9 expressing AML cell lines by transducing AML cells with a constitutive expression vector for the Cas9-mCherry fusion protein kindly provided by Marco Herold (Aubrey, Kelly et al. 2015). Then, the transduced cells (Cas9-mCherry<sup>+</sup>) were sorted using a BD Fusion BSL2 flow cytometer and expanded in culture. For the desired knockout studies, the stable AML cell line expressing Cas9-mCherry was transduced with gRNA vectors expressing either GFP, BFP or iRF670. For tracking the cells with Cas9-




sgRNA, flow cytometry analysis was used to detect double positive cells for example mCherry<sup>+</sup>iRFP670<sup>+</sup>.

### 3.4 FLOW CYTOMETRIC ANALYSIS AND SORTING

Flow cytometry is a potent tool to characterize cells based on size, granularity and fluorescence emission of a targeted antibody. The first flow cytometry analysis was developed in 1956 to detect the size of the cells and in 1980 the first clinical application of a fluorescent activated cell sorter (FACS) was used by researchers to detect lymphocyte subsets (Wilkerson 2012).

The basic principle of a flow cytometry instrument is characterization of cells in a fluid stream passing the cells through the beam of a light source (laser). Therefore, the main components of almost all flow cytometers or cell sorters are: fluidics to direct sheath buffer (phosphate-buffered saline) containing cells to the flow chamber (where the laser beams are located) by pressurized lines, optics including excitation and detection, an electronic network to convert light or fluorescent signals coming from the cells into digital signals and a computer to record digital signals (Figure 12A).



**Figure 12. Overview of flow cytometer (A) and sorter (B).** Figure reprinted with permission from the publisher (Adan, Alizada et al. 2017).

Flow cytometry is used to characterize cells based on detection of their surface-bound, cytoplasmic and nuclear antigens. Moreover, it is possible to investigate cellular components such as organelles, nuclei, DNA, RNA, cytoplasmic protein, nucleus proteins using flow cytometry.

The principle of cell sorting is an electrostatic deflection of charged droplets containing individual fluorescent cells. Flow sheath buffer containing cells is injected through a vibrating nozzle that break up the stream into droplets surrounding individual cells. Then, the droplets pass through one or more laser beams to characterize the cell population of interest and simultaneously, the tagged droplets are charged by a charging electrode. Then a platinum plate of negative charge deflects positively charged droplets and a positively charged platinum plate deflects negatively charged droplets and uncharged droplets goes into the waste container. Single or multiple parameters can be used for sorting the desired cell population and to collect them for further assays (Figure 12B).

Here flow cytometric analysis was performed with a 4-laser BD LSRFortessa.

To analyze primary childhood and adult AML cells after *ex vivo* culture, the cells were harvested and incubated with anti-mouse (Biolegend) and human (ChromPure Mouse IgG, Jackson ImmunoResearch) CD16/32 (Fc-block) antibodies for 20 minutes on ice to block unspecific receptors. Then the cells were stained for 20 minutes with anti-human CD45 to distinguish human AML cells from mouse stromal feeder cells, lineage antibodies (CD20, CD4, CD8, CD2, CD56, CD235b, CD3 and CD19) to exclude mature cells and CD34, CD38 to identify LICs. Dead cells were excluded using the Near-IR Live/Dead marker (Invitrogen). To count the absolute number of AML cells in each well from 96-well plate format, we used a high-throughput automated plate reader (BD LSRFortessa).

To detect cells with a specific fluorescent marker mixed with control cells from cell growth competition assays, cells were harvested and washed with cold PBS and thereafter stained with Near-IR Live/Dead marker in a 96-well plate in 80 µl of PBS and 2% FBS. A high-throughput automated plate reader was used to detect the absolute number of live cells.

For *in vivo* studies, and to determine the level of engrafted human AML cells in transplanted NSG-SGM3 mice, the BM cells were isolated by crushing the tibia and femur of recipient mice. The isolated BM cells were incubated in mouse and human FC blocking antibodies for 10 minutes on ice, then stained with human anti-CD45 for 20 minutes. Near-IR Live/Dead marker was used to detect live cells. To analyse the different types of mature human cell populations in NSG-SGM3 transplanted mice, the BM cells were isolated and stained with CD11b, CD14, CD56, CD19, CD33, CD16, CD3, HLA-DR and CD45.

To determine engraftment of mouse CD45.1<sup>+</sup> cells in recipient CD45.2<sup>+</sup> mice, the BM cells were isolated from tibia and femurs of recipient mice. The isolated cells were then incubated 10 minutes with mouse FC blocking antibody (anti-CD16/CD32), then 20 minutes with mouse antibodies for CD45.1 and CD45.2 to separate donor and recipient cells. To detect percentage of engrafted LSK cells, we added additional anti-mouse lineage markers (CD11b, Gr-1, CD3, CD19 NK1.1 and TER119) together with CD117 (cKit) and Sca-1. Propidium iodide (PI) (Invitrogen) was used to exclude dead cells after washing cells.

All flow cytometry data analysis was done using FlowJo Version 9.3.3 software (TreeStar).

### **3.5 CELL GROWTH ASSAYS OF AML PATIENT SAMPLES**

Long term culturing of the AML samples was carried out as previously reported (Griessinger, Anjos-Afonso et al. 2014). Briefly, mesenchymal MS-5 cells (DSMZ) were irradiated with 80 Gy, washed, and resuspended at a density of 10,000 cells/100µl of Myelocult media H5100 (StemCell Technologies Inc.). To seed feeder cells, 100 µl of cell suspension were plated in a collagen I Cellware 96-well plate (Corning), in triplicates for each condition. 2-3 days after plating feeder cells, 10,000-20,000 AML patient samples were then suspended in Myelocult media supplemented with rhIL-6, rhIL-3, rhFlt3/Flk-2 ligand, rhTPO, and rhSCF and rhG-CSF (Stemcell technology) at a concentration of 20 ng/ml, which were added to each well containing a MS-5 monolayer. The plates were kept in a box with water to increase the humidity and maintained under normoxic conditions for 3-5 weeks. The effects on cell growth (bulk or LICs) were determined by flow cytometric analysis.

### **3.6 ISOLATION AND CULTURE OF PRIMARY NORMAL CELLS**

To isolate cKit<sup>+</sup> or LSK cells, lineage mature blood cells were depleted from BM cells of the femur and tibias of C57BL/6 mice by staining cells with purified antibodies against Ter119, B220, Gr1, CD3, NK1.1, and CD11b (Biolegend) and Dynabeads (Invitrogen). Fluorochrome-conjugated anti-CD117 (cKit) and anti-Sca-1 antibodies were used to sort cKit<sup>+</sup> or LSK cells with a FACS Aria III (BD). The dead cells were excluded by PI (Invitrogen). Normal cKit<sup>+</sup>/LSK cells were cultured in SFEMII media (Stemcell technology Inc.) including; rhFlt3/Flk-2 ligand (Stemcell technology Inc.), rhTPO (Stemcell technology Inc.), rhIL-6 (R&D system), rhIL-3 (R&D systems) and rhSCF (R&D systems) at a concentration of 20 ng/ml.

For enrichment of normal human CD34<sup>+</sup> cells, Lymphoprep solution (Invitrogen) was used for the isolation of the mononuclear cell fraction from the umbilical cord blood cells (UCBs) and the CD34<sup>+</sup> cells were enriched using a CD34 magnetic activating cell sorting microbead kit (Miltenyi Biotec). Enriched CD34<sup>+</sup> cells were cultured in SFEMII media supplemented with rhIL-6, rhIL-3 (R&D systems), rhFl3/Flk-2 ligand, rhTPO and rhSCF all in a final concentration of 20 ng/ml (R&D systems) for 14 days.

### **3.7 COLONY FORMING UNIT ASSAY**

The colony forming unit (CFU) assay is an *in vitro* assay used to investigate proliferation and differentiation patterns of HSCs and progenitors by assessing their capacity to form colonies in a semi-solid medium. The morphology and number of the colonies originating from a certain number of input cells provides information about the proliferation and differentiation ability of the input cells. Moreover, serial plating of each colony illustrates preliminary information about self-renewal capacity of input HSCs.

To investigate the role of a target gene in proliferation and differentiation of HSPCs, knockdown and control groups of normal mouse (cKit<sup>+</sup>/LSK) or human BM (Lin<sup>-</sup> CD34<sup>+</sup>CD38<sup>-</sup>) were resuspended in the MethoCult semi-solid media (Stem Cell Technologies) in concentrations of 150-300 cells/ml (M3434) for mouse and 200-400 cells/ml (M4435) for human. After vortexing the mixture vigorously, 1 ml of the cell mixture was seeded in a 1 cm<sup>2</sup> dish with a 16g gauge blunt-end needle. The colonies were counted and characterized according their morphology using an inverted microscope at 10–12 days (mouse) or 12–14 days (human) after plating. A cluster of more than 50 cells was defined as one colony.

### **3.8 AML MOUSE MODELS AND TRANSPLANTATION STUDIES**

For the *in vivo* part of studies I, II, III and V, all the mice were kept in a specific pathogen-free animal facility at Karolinska Institutet, Huddinge, Sweden. The C57BL/6 wild type mice and the NOD Scid IL2Rgnull-3-SCF/GM/IL3 (NSG-SGM3) mice were purchased from The Jackson Laboratory. All the transplanted mice were monitored daily for symptoms of leukemogenesis and disease progression was investigated using complete blood tests.

The mouse *MLL-AF9* AML cells were generated according to a previous study (Somervaille and Cleary 2006). cKit<sup>+</sup> cells were sorted from the BM of CD45.1<sup>+</sup> C57BL/6 mice and they were transduced with a retroviral vector expressing the MLL-AF9 fusion oncogene. The

transduced cells were kept in *in vitro* culture and were serially re-plated in semi-solid methylcellulose medium containing 20 ng/ml rmSCF, 10 ng/ml rhIL-6, 10 ng/ml rmGM-CSF, 10 ng/ml rmIL-3 and G418 (0.75 mg/ml) to enrich immortalized cells. Single immortalized clones were expanded and transplanted via the tail vein into sub-lethally irradiated (600 cGy) CD45.2<sup>+</sup> C57BL/6 wild type mice. Once transplanted mice showed symptoms of leukemogenesis and disease progression, MLL-AF9 AML cells were harvested from the BM of euthanized primary AML mice, sorted by flow cytometry and expanded in RPMI medium in the presence of IL3 (10 ng/ml) and 10% FBS. The sorted mouse AML cells were used for further *in vitro* or *in vivo* experiments.

For the Kaplan-Meier survival analysis of the AML mouse model, 250,000 mouse *MLL-AF9* AML cells after transduction and selection with puromycin (2 µg/ml) were intravenously transplanted into wild type CD45.2 C57BL/6 mice aged 6-8 weeks. After AML onset, the mice were euthanized.

For *in vivo* experiments with primary human cells including normal hematopoietic cells and AML patient samples, we used sub-lethally irradiated (220 cGy) humanized NSG-SGM3 mice aged 6-10 weeks. The primary cells (normal cells: 25,000-50,000 and AML patient: 100,000-500,000) samples were transplanted via intra-femoral injection. For detection of engrafted cells, the recipient mice were euthanized and the BM cells were collected and analyzed by flow cytometry.

### 3.9 RNA SEQUENCING

RNA sequencing is a powerful technology to investigate the RNA expression profile of cells. Briefly, extracted RNA, either total or poly-A<sup>+</sup> mRNA is converted to cDNA fragments and adaptors are attached in one or two ends of the fragments. The library fragments can be sequenced with high throughput from one end (single-end sequencing) or both ends (pair-end sequencing). The length of the sequenced reads is around 30-400 bp depending on the sequencing technology used. To analyze the reads after sequencing, they are either aligned to a reference genome or reference transcript, or they can be assembled *de novo* without the genome sequence. Then mapped data are normalized and bioinformatic analyses such as expression level, expression of isoforms and whole transcriptome variation and statistical analyses can be performed.

For RNA sequencing in studies I, II, III, and V, total RNA was extracted from cells using the RNeasy Micro Kit (Qiagen) 72 hours after transduction of the cells by sorting or selection

with puromycin. TotalScript™ RNA-seq kit (Epicentre, Madison, WI) was used to prepare strand specific pair-end RNA libraries according to the manufacturer's instructions. Libraries were pair-end sequenced using the Illumina platform HiSeq2000 or Nextseq500. RNA-seq reads were mapped to the Ensembl Homo sapiens GRCh38 reference genome using the STAR aligner. Gene assignment was performed using feature Counts. Normalization and the sample group comparison was performed using DESeq2.

## 4 RESULTS AND DISCUSSION

### 4.1 RESULTS

#### 4.1.1 Study I

In this study, we aimed to identify novel target genes that are selective and essential for maintenance of AML. For this purpose, we performed shRNA based screens of the human AML cell lines NOMO-1 and a mouse AML cell line. All cell lines were characterized by MLL-AF9 translocation. To filter out targets with a strong effect on normal hematopoietic cells we used non-transformed mouse Factor-Dependent Continuous Paterson Laboratories (FDCP mix) cells as control cells in the screens (Spooncer, Heyworth et al. 1986). The AML cells were transduced with lentiviral virus from an RNAi library consisting 27,000 shRNAs targeting 5,400 putative disease-associated and drug targets (Cellecta Inc.). The transduced cells were harvested at the initial time point (T0) and after ten cells divisions (T10) to sequence the PCR-amplified unique barcodes for individual shRNAs. By comparing the ratio of each individual barcode at two time points, we identified 1,082 target genes with at least a five-fold higher inhibitory effect on mouse AML cell growth compared to FDCP-mix cells. Among the 1,082 targets, 34 were common between THP-1 and NOMO-1 cell lines as shown by at least a five-fold effect in the screens.

Among 34 target genes, multiple shRNAs were scored against chromodomain helicase DNA-binding protein 4 (CHD4) which had a strong growth inhibitory effect on AML cell lines. To validate whether CHD4 is required for AML cell growth, we performed an *in vitro* competition assay by mixing shRNA expressing RFP and targeting *Chd4* with control shRNA tagged with a GFP fluorescent marker. Knockdown of *Chd4* with two independent shRNAs caused a strong blockade of leukemic cells in THP-1, NOMO-1 and mouse AML cells. We then tested the effect of *CHD4* knockdown in non-MLL AML cells lines by using HL-60, K562 and NB-4 AML cell lines. The result from this *in vitro* growth assay revealed that *CHD4* is also crucial for growth of non-MLL AML cells. We used CRISPR/Cas9 technology to target *CHD4* on two human AML cell lines (THP-1, MV4-11) to avoid possible off-target effects commonly caused by shRNA, and confirmed that *CHD4* expression is crucial for propagation of AML cells.

To investigate whether CHD4 plays role in AML progression, we transplanted MLL-AF9 AML cells into recipient congenic mice. The mice which received *Chd4* knockdown AML cells survived significantly longer than the control group transplanted with AML cells

carrying Sc control shRNA. To check whether *CHD4* is important for normal hematopoietic cells, we first tested the effect of knockdown on normal mouse HSPCs in a growth assay format. The flow cytometric analysis demonstrated a minor difference between cells transduced with *Chd4* shRNA and control cells. Moreover, a similar result was seen when testing *CHD4* shRNA in human CD34 positive UCBs. Together, these results indicate that *CHD4* plays an important role in the growth of leukemic cells *in vitro* but not in normal primary hematopoietic cells.

Next, we tested knockdown of *CHD4* shRNA in primary childhood patient samples *ex vivo* by culturing transduced patient samples either with control shRNA or *CHD4* shRNA on MS5 stromal feeder cells. Flow cytometry analysis demonstrated that the expansion of cells transduced with *CHD4* shRNA was significantly inhibited compared to control cells, whilst the absolute number of LICs were decreased in *CHD4* knockdown cells. These results suggest that *CHD4* inhibition prevents expansion of bulk leukemic cells as well as LICs. Interestingly, using a humanized NSG-SGM3 mouse model to transplant childhood patient samples transduced with either *CHD4* shRNA or Sc control showed that primary childhood AML cells transduced with *CHD4* shRNA had significantly lower levels of leukemic engraftment in recipient mice compared to the Sc control group.

Cell cycle assay analysis of MLL rearranged AML (THP-1 and MV4-11) and non-MLL AML (AML-193 and Kasumi-1) human cell lines revealed that inhibition of *CHD4* caused an accumulation of cells in G0 phase of cell cycle. Additionally, *CHD4* inhibition reduced the number of cells in G1 phase but did not significantly affect the number of cells in S and G2/M phases compared to control cells. Contrastingly, a minor effect on apoptotic cells was observed using *CHD4* shRNA.

Next, to determine the RNA expression pattern induced by *CHD4* knockdown, we performed RNA sequencing analysis of THP-1 AML cells transduced with *CHD4* shRNA. Gene set enrichment analysis (GSEA) of RNA-seq data showed that the genes most significantly correlated with *MYC* and *E2F* targets. Moreover, knocking down of *CHD4* induced downregulation of *MYC* and genes involved in G1/S cell cycle such as D1, D2, E1, E2F1 and E2F2.

In this study, we found that the shRNAs targeting *CHD4* identified from our screens targeted AML cells without a dramatic effect on normal hematopoietic cells, making *CHD4* a potential therapeutic option for AML. We showed that *CHD4* knockdown reduced the frequency of bulk AML cells as well as LICs in primary childhood AML samples. Additionally, we determined that *CHD4* is essential for AML progression and development



by testing the effect of knockdown in a congenic AML mouse model and transplanting patient samples into NSG-SGM3 mice. These data revealed an important role for *CHD4* in childhood AML.

#### 4.1.2 Study II

In this study, our goal was to identify novel TFs essential for AML cell expansion. We screened two human cell lines (THP-1 and NOMO-1) and mouse AML cells with a lentiviral shRNA library targeting around 5,000 genes within signaling pathways (Cellecta Inc.). The transduced cells were harvested at two time points; initial and after 10 cell divisions. Each individual shRNA vector contained a unique barcode which was amplified from genomic DNA and tracked by NGS. We found 648 target genes which caused at least a five-fold reduction in AML cell growth overlapping between THP-1, NOMO-1 and mouse AML cells. Among the target genes, 38 of them were TFs and 24 of the 38 were known to be important in AML such as *MEIS1*, *HEMS1* and *MYC*. To avoid strong effects on normal cells, we prioritized expression levels of TFs in myeloid leukemia cells and normal hematopoietic cells as listed in the BloodSpot data base. *GTF2IRD1* is one such TF that was highly expressed in leukemic cells.

To validate *GTF2IRD1*, we first tested mouse AML cells by performing growth assays using AML cells transduced with shRNAs targeting *Gtf2ird1*, mixed with cells transduced with Sc control shRNA. FACS analysis showed *Gtf2ird1* knockdown significantly impaired AML growth. We found a similar result in five human AML cell lines; THP-1, NOMO-1, HL-60, K-562, and NB-4. Then we investigated the importance of *Gtf2ird1* in AML development *in vivo* using a congenic AML mouse model. The mice transplanted with *Gtf2ird1* knockdown AML cells survived longer compared to the recipient mice transplanted with AML cells transduced with Sc control shRNA.

Our next question was if *GTF2IRD1* blocks the growth of normal hematopoietic cells. To answer this question, we first performed *in vitro* growth assays on mouse HSPCs. FACS data showed no significant difference between cells transduced with *Gtf2ird1* shRNA and control cells. Furthermore, the CFU assay confirmed *Gtf2ird1* is not essential for expansion and differentiation of mouse HSPCs *in vitro*. Transplantation of depleted *Gtf2ird1* HSPCs into lethally irradiated mice and monitoring the frequency of engraftment demonstrated that knockdown of *Gtf2ird1* did not strongly affect the reconstitution of HSPCs cells *in vivo*.

To explore the effect of *GTF2IRD1* inhibition on survival and expansion in normal human hematopoietic cells, we performed a growth assay on CD34<sup>+</sup> cells enriched from UCBs

transduced with *GTF2IRD1* shRNA or Sc shRNA. Similar to the results observed with mouse HPSCs, we again saw that *GTF2IRD1* is not crucial for expansion of human hematopoietic cells. Long-term culturing of transduced CD34<sup>+</sup> UCBs cells also revealed *GTF2IRD1* inhibition did not influence normal cell growth. Transplantation of CD34<sup>+</sup> UCBs into NSG-SGM3 mice and FACS analysis of engrafted cells after 8 weeks demonstrated no significant differences in engraftment between the mice receiving cells with *GTF2IRD1* knockdown compared to the Sc control group. Altogether, data from mouse and human HSPCs suggest that the *GTF2IRD1* TF has a selective role in the expansion and development of AML but is not important in normal hematopoiesis.

Additionally, we knocked-out *GTF2IRD1* in THP-1 and MV4-11 cells using CRISPR/Cas9 technology and performed an *in vitro* competition assay. By comparing cells with gRNA targeting *GTF2IRD1* to cells transduced with control gRNA, we observed a strong reduction in *GTF2IRD1* knockout cells which confirmed our result as previously seen in AML cells using shRNA.

To investigate the role of *GTF2IRD1* in primary childhood and adult AML patients, we took advantage of *ex vivo* culturing of AML samples on stromal feeder cells. Transduction of patient samples with *GTF2IRD1* shRNA caused a significant reduction of bulk AML (CD45<sup>+</sup>) cells as well as LICs (CD45<sup>+</sup>Lin<sup>-</sup>CD34<sup>+</sup>Cd38<sup>-</sup>). Additionally, intrafemoral transplantation of transduced cells into NSG-SGM3 humanized mouse cells showed *GTF2IRD1* knockdown decreased the percentage of engrafted adult and childhood AML samples. Altogether, these data confirm that expression of *GTF2IRD1* is crucial for the survival and maintenance of adult and childhood primary AML cells.

Cell cycle and apoptosis analysis of mouse AML cells revealed a significant increase in the number of cells in the G0 cell cycle phase and reduction in G1, S and G2M phases caused by knocking down of *Gtf2ird1* compared to control cells. In contrast, the percentage of apoptotic cells at the same time point was low, around 2-3%, suggesting that *Gtf2ird1* knockdown induces an accumulation of cells in the G0 phase of the cell cycle rather than inducing apoptosis.

To find out the role of *GTF2IRD1* in gene regulation, we used a heterologous GAL4 reporter system (Hansen, Bracken et al. 2008). In this system, the luciferase reporter gene is regulated by a thymidine kinase promoter with five GAL4 binding sites, therefore suppressor and activation function of a TF fused with GAL4 can be measured by luciferin expression level. Tetracycline induction of *GTF2IRD1*-GAL4 expression suppressed luciferase expression levels, suggesting that *GTF2IRD1* TF acts as a transcriptional suppressor in AML cells.

RNA sequencing of mouse AML cells with *Gtf2ird1* knockdown compared to control cells revealed that *Gtf2ird1* knockdown mRNA profiles were enriched in various cellular pathways, including myeloid leucocytes activation, lipid biosynthesis and cell-cell adhesion. Furthermore, GSEA showed knockdown of *Gtf2ird1* negatively correlated with *MYC* targets data sets (FDR q value=0.0 and NES = -1.7) and KRAS signaling (FDR q value = 0.0, NES -1.6).

### 4.1.3 Study III

Here we focused on the role of euchromatin histone methyltransferases 1 (EHMT1) in AML. EHMT1 and EHMT2 are enzyme homologs which specifically methylate the lysine 9 (K9) residue in histone 3 (H3), acting as transcription repressors. Several studies have shown the function of EHMT2 in AML (Lehnertz, Pabst et al. 2014), but the role of EHMT1 in AML is not currently clear.

To first investigate the importance of *Ehmt1* in AML, we knock downed *Ehmt1* using shRNA in mouse MLL-AF9 AML cells and analyzed the growth of AML cells in competition with control cells. Monitoring cells via flow cytometry analysis for two weeks showed that knockdown of *Ehmt1* significantly reduced the frequency of AML cells *in vitro*. To further analyze the role of *Ehmt1* in mouse AML cells, we transplanted AML cells into lethally irradiated recipient mice with and without knockdown of *Ehmt1*. Interestingly, transplanted cells containing one of the independent shRNAs targeting *Ehmt1* prolong the survival of mice in comparison to mice transplanted with control cells, which died 18-21 days after transplantation. These data suggest that expression of *Ehmt1* is necessary for the growth and progression of mouse AML cells.

Next, we studied the importance of *EHMT1* in human AML cells lines by transducing them with shRNA against *EHMT1*. Growth assay analysis of five AML cells lines with MLL-AF9 translocation (THP-1 and NOMO-1) and without MLL-AF9 alteration (HL-60, K-562 and NB-4) revealed that the essential role of *EHMT1* in mouse AML cell expansion was conserved in human AML cells as well. Moreover, suppression of *EHMT1* activity with the small inhibitor molecule; BIX-01294 in human KL-60 and THP-1 AML cells showed similar results in line with *EHMT1* shRNA; namely that inhibition of *EHMT1* significantly impairs the growth of AML cells.

To confirm the results observed using shRNA, we used CRISPR/Cas9 to knockout *EHMT1* in human AML cell lines. We transduced THP-1 and MV4-11 cell lines with stable expression of Cas9-mCherry using a gRNA targeting the *EHMT1* gene. Targeting *EHMT1*

with gRNA caused a significant reduction in AML cell growth, confirming our previous result regarding the important role of *EHMT1* in AML cell growth.

To find out whether *EHMT1* is important for normal hematopoietic cell propagation we tested the effect of *EHMT1* knockdown on mouse and human HSPCs. In both cases we observed a reduction in cell growth, but this reduction in cell number was not as strong as seen in AML cells. To further investigate, we performed CFU assays for normal mouse and human HPSCs with and without *EHMT1* knockdown. The results revealed *EHMT1* suppression caused a reduction in colony number. Together, these data suggest that AML cell growth is more dependent on *EHMT1* compared to normal HSPCs.

Next, to demonstrate the role of *EHMT1* in primary AML adult and childhood patient samples, we co-cultured transduced cells with either *EHMT1* or Sc shRNA with MS5 stromal feeder cells (Griessinger, Anjos-Afonso et al. 2014). Flow cytometry analysis to read out the frequency of bulk AML cells ( $CD45^+$ ) and LIC ( $Lin^-CD34^+CD38^-$ ) showed that *EHMT1* knockdown AML cells expanded less compared to control cells. To further investigate the role of *EHMT1* in the expansion of primary AML cells, we transplanted *EHMT1* knockdown cells into a NSG-SGM3 mouse model and monitored the engrafted cells by flow cytometry up to eight weeks post transplantation. This was in comparison to control mice receiving AML cells transduced with Sc shRNA. Engraftment of primary patient samples was significantly lower in the group with *EHMT1* knockdown. Therefore, these data showed *EHMT1* also plays an important role in the growth of primary AML samples in *in vitro* and *in vivo* conditions.

*EHMT2* is another member of the EHMT family, described as a heterodimer to *EHMT1* (Tachibana, Ueda et al. 2005), with important roles in AML previously reported (Lehnertz, Pabst et al. 2014). To address the question if *EHMT1* and *EHMT2* have an overlapping role in AML, we designed gRNAs against *EHMT1* or *EHMT2* with different fluorescent reporter markers, enabling us to track them in co-culture conditions and to detect cells carrying both gRNAs. Flow cytometry analysis of the competition assay revealed that single knockout of *EHMT1* or *EHMT2* reduced AML growth equally and interestingly, double knockout of *EHMT1* and *EHMT2* did not show an additive effect on AML cell growth. These data suggest that *EHMT1* and *EHMT2* have a cooperative function in the growth of AML cells.

To understand the cellular mechanism of *EHMT1* and *EHMT2* in AML cells, we performed cell cycle analysis on mouse MLL-AF9 AML cells with and without knockdown for either *Ehmt1* or *Ehmt2*. Interestingly, we observed both *Ehmt1* and *Ehmt2* knockdown caused a significant increase in the proportion of cells in the G0 phase of the cell cycle and

simultaneous reduction of cells in S and G2M phases. However, we did not observe a significant difference in the number of apoptotic cells, suggesting both *EHMT1* and *EHMT2* inhibition induce cell cycle arrest in AML cells in G0.

To uncover the molecular mechanism of *Ehmt1* in the regulation of AML cell growth, we RNA sequenced mouse AML cells with *Ehmt1* knockdown. Additionally, to detect similarities and differences in the RNA expression profiles between *Ehmt1* and *Ehmt2*, we performed RNA-seq analysis of *Ehmt2* knockdown AML cells. The mRNA profile of *Ehmt1* showed a significant correlation to the *Ehmt2* mRNA profile and since both enzymes are transcriptional suppressors, they unsurprisingly showed a significant overlap in the upregulated genes. Moreover, GO-term analysis of upregulated genes revealed common biological processes associated to cytokine signaling, inflammatory response and cell differentiation for *Ehmt1* and *Ehmt2*. These RNA-seq data suggest although *Ehmt1* and *Ehmt2* share several biological processes, they may still have further independent roles as well.

#### **4.1.4 Study IV**

In this study, we contributed to Nikolas Herold's study to investigate the effect of SAMHD1 inhibitor Vpx, in the treatment of patient AML samples with the cytostatic deoxycytidine analog cytarabine (ara-C). ara-C is the most common drug for AML treatment and resistance to ara-C is the main cause of CR failure and relapse.

SAMHD1 was identified as a risk factor in cohorts of both childhood and adult AML patients who received ara-C treatment. Using simian immunodeficiency virus (SIV) protein Vpx to transiently reduce SAMHD1 expression, AML cells demonstrated a markedly increased sensitivity to ara-C-induced cytotoxicity. Moreover, disruption of *SAMHD1* using CRISPR/Cas9 technology showed a similar effect in increasing sensitivity of AML cells to ara-C.

Xenotransplantation of THP-1 cells with or without expression of *SAMDHI* into athymic nude immunodeficient NMRI nu/nu mice and treatment of the animals with ara-C revealed that SAMHD1<sup>-</sup> tumors responded to the treatment compared to SAMHD1<sup>+</sup> tumors.

Next, we treated adult and childhood AML patient samples under *ex vivo* conditions with virus-like particles (VLPs) either with or without Vpx. Flow cytometric results revealed that reduction of SAMHD1 protein levels in patient AML samples increased sensitivity of cells

to ara-C. We also showed that Vpx increased the number of apoptotic cells compared to cells treated with empty VLP.

Thus, a low expression level of SAMHD1 is associated with sensitivity of leukemic cell to ara-C, suggesting that the targeting of SAMHD1 could be used as a potential therapeutic strategy to increase ara-C efficacy in AML patients.

#### 4.1.5 Study V

Epigenetic regulators play an important role in hematopoiesis. In particular, the self-renewal capacity and differentiation abilities of HSCs, acting as the main source of all mature blood cells are regulated with epigenetic regulators. In this study, we investigated the role of *NAP1L3*; nucleosome assembly proteins in self-renewing and differentiation of HSCs *in vitro* and *in vivo* and signaling pathways regulated by *NAP1L3* in hematopoiesis. *NAP1L3* has been reported as one of the 36 transcriptional regulatory genes expressed predominantly in HSCs, suggesting a potential role in HSC regulation (Riddell, Gazit et al. 2014). To detect the expression level of *Nap1l3* in different populations of mouse hematopoietic and progenitor cells, we measured the *Nap1l3* mRNA level in seven FACS sorted populations; HSC (Lin<sup>-</sup>Sca1<sup>+</sup>cKit<sup>+</sup>CD105<sup>+</sup>CD150<sup>+</sup>), multi-potent progenitors (MPP; LSK<sup>+</sup>CD105<sup>+</sup>CD150<sup>+</sup>), lymphoid-primed multipotent progenitors (LMPP; LSK<sup>+</sup>Flk2<sup>high+</sup>), common lymphoid progenitors (CLP; Lin<sup>-</sup>IL7Ra<sup>+</sup>flk2<sup>+</sup>), pre-granulocyte-macrophage progenitors (pre-GM; LSK<sup>-</sup>CD41<sup>-</sup>CD150<sup>-</sup>CD105<sup>-</sup>), granulocyte-monocyte progenitors (GMP; LSK<sup>-</sup>CD41<sup>-</sup>CD150<sup>-</sup>FcgR<sup>+</sup>), and erythrocyte progenitors (pre-CFU E; LSK<sup>-</sup>CD41<sup>-</sup>CD105<sup>+</sup>). Consistent with previous studies, qPCR analysis showed that *Nap1l3* was exclusively expressed in the HSC fraction.

To explore the importance of *Nap1l3* in HSPCs proliferation and differentiation, shRNA-based knockdown was used to study loss of function of *Nap1l3*. Knockdown of *Nap1l3* in FACS sorted mouse Lin<sup>-</sup>Sca1<sup>+</sup>cKit<sup>+</sup> (LSK) cells caused a significant reduction of total CFUs, as well as mixed myelo-erythroid CFUs (CFU-GEM) and granulocyte/macrophage CFUs (CFU-GM) numbers compared to the LSK cells transduced with negative scramble control shRNA. These data suggest that *Nap1l3* suppression impairs the proliferation and survival of mouse HSPCs *in vitro*.

To avoid potential off-target effects associated with shRNAs, we designed gRNAs to target *Nap1l3* via CRISPR/Cas9 technology. LSK cells were sorted from transgenic mice overexpressing Cas9 nuclease and then transduced with gRNA vectors against *Nap1l3*.

Consistent with shRNA-mediated knockdown, knockout of *Nap1l3* with CRISPR/Cas9 caused a significant reduction in the CFU colony number compare to control cells.

Next, we used exogenous expression vectors to induce expression of *Nap1l3* in mouse HSPCs. Constitutive overexpression of *Nap1l3* caused a significant reduction of the total number of CFUs, CFU-GM and CFU-GEM colonies.

To investigate this further, we transplanted sorted (ckit<sup>+</sup>) HSPCs transduced with *Nap1l3* shRNA or Sc shRNA into congenic lethally irradiated mice. We then monitored the engrafted cells (CD45.1<sup>+</sup>) in the BM of recipient mice 2, 5, 8 and 16 weeks post-transplantation. Flow cytometric analysis revealed knockdown of *Nap1l3* caused a significant reduction in engraftment. Moreover, by deep analysis of the BM population, we showed a distinct reduction in LSK cells mediated by *Nap1l3* downregulation. These data suggest that *Nap1l3* plays an important role in the survival and reconstitution of HSC in both the short- and long-term *in vivo*. Furthermore, flow cytometric analysis of mature blood cells showed a decrease in the frequency of myeloid cells (CD11b<sup>+</sup>), granulocytes (Gr-1<sup>+</sup>) and, contrastingly, an increase B cells (CD19<sup>+</sup>) in *Nap1l3* knockdown cells. Altogether, these data show *Nap1l3* may have a role in differentiation regulation of HSCs.

To dig in to the importance role of *NAPIL3* in human HSCs, we transduced enriched HSCs (Lin<sup>-</sup>CD34<sup>+</sup>CD38<sup>-</sup>) from UCB cells with two shRNAs against *NAPIL3* and Sc control. Proliferation analysis of these cells after 48 hours showed that *NAPIL3* knockdown impaired proliferation and reduced the number of mature cells (Lin<sup>+</sup>) compared to control cells, suggesting that *NAPIL3* is required for proliferation and differentiation of HSC *in vitro*. Culturing UCB HSCs with *NAPIL3* inhibition on stromal feeder cells (SL/SL and M2-10B4) for three weeks showed a significant reduction in CD45<sup>+</sup>, Lin<sup>-</sup>CD45<sup>+</sup>, and UCB HSCs. Also, CFU assays on normal human UCB HSCs demonstrated that knockdown of *NAPIL3* significantly reduced the number of colonies as well as burst-forming unit erythroid cells (CFU-E/BFU-E), macrophages (CFU-M), granulocytes/macrophages (CFU-G/GM), and mixed myelo-erythroid cells (CFU-GEM). To study the effect of *NAPIL3* in HSC self-renewal, we performed a serial plating assay. In doing so, we observed a reduction of colony numbers in the first, second and third plating which suggests *NALP13* may also be important for HSC self-renewal.

To study the cellular mechanisms controlled by *NAPIL3*, we performed cell cycle analysis and apoptosis assays. Cell cycle analysis of UCB HSCs with *NAPIL3* knockdown compared to control cells revealed an accumulation of cells in the G0 phase and a reduction of cells in

G1. Apoptosis assays of cells showed an increased population in early apoptosis. Altogether, knockdown of *NAPIL3* induced cell cycle arrest of HSCs in G0 and triggered apoptosis.

Next, we wanted to see the role of *NAPIL3* *in vivo*, so we transplanted CD34<sup>+</sup> HSPC UCBs transduced with *NAPIL3* or Sc shRNA into humanized NSG-SGM3 mice. Flow cytometric analysis of BM cells in recipient mice 8 weeks post-transplantation showed that *NAPIL3* knockdown dramatically reduced engraftment percentage. FACS analysis of human mature cells in recipient mice also showed an increase in the percentage of myeloid cells (monocytes, dendritic like cells, plasmacytoid dendritic cells) compared to the control group. Therefore, *NAPIL3* is likely to play an important role during HSC self-renewal and differentiation.

To uncover the mRNA expression profile induced by *NAPIL3* knockdown, we performed RNA sequencing of sorted UCB HSCs transduced with *NAPIL3* compared to cells transduced with control Sc shRNA. GSEA analysis of Biocarta gene set pathways revealed that the mRNA profile of cells with *NAPIL3* knockdown negatively correlated to cell cycle regulation, chromosome function, recombination and replication. We also observed *NAPIL3* knockdown induced upregulation of the *HOXA* cluster in RNA-seq data, which we confirmed by qPCR for all *HOXA* clusters except *HOXA4*. Flow cytometry also revealed that protein levels of HOXA3 and HOXA5 were upregulated in the presence of *NAPIL3* shRNA. In conclusion, these data show that downregulation of *NAPIL3* in UCB HSCs induces gene expression signatures associated with cell cycle progression and *HOXA* gene expression.

## 4.2 DISCUSSION

To design new drugs to eradicate cancer cells, it is very important to selectively target the cancer cells without eliminating normal cells. In studies I and II, we found that *CHD4* and *GTF2IRD1* play critical roles in leukemic cell growth in various AML cell lines with different mutational backgrounds and in AML development in mouse models. Interestingly, they are not pivotal for the normal function of hematopoietic cells.

In **study I**, we identified the chromatin remodeler *CHD4* from shRNA screens of mouse and human AML cell lines and non-transformed FDCP cells. Our results illustrate a novel and crucial role of *CHD4* in the maintenance and progression of childhood AML by testing the patient samples in an *ex-vivo* co-culture system and a patient-derived xenograft mouse model. However, *CHD4* knockdown did not show a significant effect on normal mouse and human hematopoietic cell growth. Our findings do emphasize a vital role for *CHD4* in the expansion of primary childhood AML and interestingly in the maintenance of LICs. Using immune



phenotyping of surface markers to identify LICs, we found that expression of *CHD4* is important for them, which is in line with other cancer types (Chudnovsky, Kim et al. 2014, Nio, Yamashita et al. 2015). Knockdown of *CHD4* revealed an important role of this epigenetic factor in the growth of AML by arresting cells in G0. We showed *CHD4* inhibition causes cell cycle arrest in G0, and our results are supported by other findings recognizing the role of *CHD4* in the cell cycle (O'Shaughnessy and Hendrich 2013, Polo, Kaidi et al. 2010). To illustrate how *CHD4* affects the cell cycle, our RNA-seq data revealed a significant downregulation of *MYC* and its target genes under *CHD4* knockdown conditions, which suggest that the *MYC* gene may be a downstream target of *CHD4* which causes cell arrest in G0. To this end, our findings illustrate *CHD4* as a potential therapeutic target, which can be used to selectively eliminate childhood AML cells and to particularly diminish LICs.

In **study II**, we found a novel TF, *GTF2IRD1*, from large scale shRNA screens and we demonstrated that both knockdown and knockout of *GTF2IRD1* selectively blocked mouse and human AML growth. We found a novel role of TF *GTF2IRD1* in maintenance and growth of both childhood and adult AML. Although overexpression of *GTF2IRD1* was reported in AML cells compared to normal hematopoietic cells, our results revealed for the first time the functional role of *GTF2IRD1* in leukemic cells as well as in LICs growth in adult and childhood AML patient samples (Bagger, Sasivarevic et al. 2016). Though more investigations are needed to uncover the molecular mechanism of *GTF2IRD1* in the growth of leukemic cells, our results suggest a selective inhibitory role in AML cell expansion but not in normal hematopoietic cells. Additionally, *GTF2IRD1* knockdown caused an accumulation of AML cells in G0 and downregulation of *MYC* targets and *KRAS* signaling. Despite the transcriptional repressor function of *GTF2IRD1*, RNA-seq data from AML cells with *Gtf2ird1* knockdown revealed a strong downregulation of *MYC* targets at the mRNA level, which might correlate with a secondary side effect of blocking cells in G0 upon *Gtf2ird1* knockdown rather than a direct regulatory role of *MYC* target genes. Recently, it has been possible to target TFs in new therapies for cancer cells by modifying their expression or degradation at the mRNA level, by inhibiting protein-protein interactions, or by direct targeting TF and block binding the TF to DNA (Lambert, Jambon et al. 2018). In particular, there are several efforts to develop new strategies to target different TFs in AML including *C/EBPα*, *PU.1*, *RUNX1*, *RUNX2*, *RUNX3*, *p53*, *c-MYC*, *CREB* and *STAT3* (Takei and Kobayashi 2018). Most importantly, we have identified a novel role of *GTF2IRD1* in AML cell growth and maintenance, suggesting a new TF as a promising therapeutic candidate to selectively target AML cells.

The aim of **study III** was to investigate the role of histone methyltransferase *EHMT1* in AML and to establish a connection to its homolog *EHMT2*. We showed that downregulation of *EHMT1* had a negative effect on maintenance of AML cells in both *in vitro* and *in vivo*. Knockdown of *EHMT1* reduced the number of bulk and LIC cells in primary AML patient samples. Our results suggest that expression of *Ehmt1*, similar to *Ehmt2*, is vital for proliferation of AML cell lines as well as patient-derived AML samples. Simultaneous knockout of *EHMT1* and *EHMT2* in AML cells did not show an additive effect on cell growth compared to single knockout, suggesting that *EHMT1* and *EHMT2* functionally overlap in the regulation of AML growth. Our studies also revealed that both *Ehmt1* and *Ehmt2* knockdown arrests cells in G0 of the cell cycle. Furthermore, our RNA-seq data showed significant overlap in upregulated genes between *Ehmt1* knockdown AML samples and *Ehmt2* knockdown samples. This is consistent with a previous study showing that *Ehmt1* and *Ehmt2* predominantly form a heterodimer complex in the nucleus and are both cooperatively important for H3K9 methylation as an inactivating epigenetic marker (Tachibana, Ueda et al. 2005). Besides that, our data also revealed several upregulated genes and pathways that are not shared between the two groups of knockdown cells, suggesting *Ehmt1* and *Ehmt2* might play independent roles in AML cells in addition to their cooperative role. Since both epigenetic enzymes are associated with transcriptional inactivation, downregulated genes most likely represent secondary effects of *Ehmt1* or *Ehmt2* knockdown. Further experiments need to be done to clarify the functional relationship between the two epigenetic enzymes and promoter regions they bind to together or individually, but it is possible to find homodimer complexes of *Ehmt1* and *Ehmt2* with specific regulatory sites in AML cells. Also, more investigations are clearly needed to identify which of the two H3K9me1/2 methyltransferases shows less pronounced effects on normal hematopoietic cells. Based on the results, studies can be performed to design a specific inhibitor that selectively blocks either EHMT1 or EHMT2 as therapeutic candidate in AML.

In **study IV**, resistance to ara-C and relapse after treatment are the main obstacles to achieve long term CR in AML patients. Therefore, finding a prognostic marker to predict chemotherapy response will be valuable for AML patients. Here, the data suggested that there is a significant correlation between high level expression of *SAMHD1* and low sensitivity to ara-C in AML patient samples, suggesting the correlation between SAMHD1 expression and prediction of ara-C treatment response. The increased sensitivity to ara-C in the absence of SAMHD1 could be because of elevated levels of intracellular ara-CTP, which cause increased incorporation of ara-C into DNA. However, there is no significant difference in SAMHD1 levels between AML patients with CR and patients who did not achieve CR. One

explanation could be clonal selection of ara-C resistant sub-clones with higher levels of SAMHD1 after therapy that might induce relapse. Moreover, the probable tumor suppressor function of SAMHD1 might increase the survival advantage of sub-clones with low SAMHD1 expression level over time (Clifford, Louis et al. 2014). Importantly, our results suggest that expression of SAMHD1 may be a limiting factor for ara-C therapies and evaluation of the level of SAMHD1 could be helpful to avoid toxicity related to increasing doses of ara-C. Further efforts are needed to explore strategies to inhibit SAMHD1 in combination with ara-C treatment, but these data suggest that leukemic cells can be sensitized to ara-C with inhibition of SAMHD1.

In **study V**, our investigations recognized a novel role of the histone chaperone *NAPIL3* in HSCs. Loss of function of the highly expressed *NAPIL3* in HSPCs significantly impaired survival, proliferation, reconstitution and differentiation of HSCs *in vitro* and in a transplanted mouse model. These data suggest the transcriptional regulatory role of *NAPIL3* on genes that are associated with cell cycle and differentiation processes. While the role of Nap1 family proteins in cell cycle regulation and during mitosis has been confirmed, we showed for the first time that a specific member of the Nap1 family, *NAPIL3*, is required for regulation of cell cycle in normal hematopoietic cells and its inhibition causes cell arrest in G0 (Altman and Kellogg 1997, Grande, Lambea et al. 2008). Our data is in line with previous data, which showed that NAP proteins can bind to core histones and physically interact with p300 coactivators to regulate target genes including p53 and E2F (Shikama, Chan et al. 2000). Knockdown of *NAPIL3* in human hematopoietic cells caused an abnormal proportion of mature cells in a xeno-graft mouse model, which might suggest a regulatory role of *NAPIL3* in the epigenetic landscape of HSCs during differentiation to produce all lineages of mature blood cells. Furthermore, *NAPIL3* knockdown decreased the reconstitution and self-renewal capacity of HSCs. Although our RNA-seq data from human HSPCs with *NAPIL3* knockdown did not show significant association with hematopoietic differentiation pathways, we showed that five *HOXA* genes were significantly upregulated upon *NAPIL3* suppression at the mRNA level and two of three investigated at protein level. Since the roles of *HOXA* genes in self-renewing and differentiation of HSCs are well known, the correlation between *NAPIL3* knockdown and upregulation of *HOXA* genes may give a clue for the role of *NAPIL3* in HSC self-renewal and differentiation (Lebert-Ghali, Fournier et al. 2016).



## 5 CONCLUDING REMARKS

In order to improve quality of life and increase survival of AML patients, it is vital to increase our knowledge regarding molecular mechanisms that drive the disease and relapse.

New advanced techniques including extensive NGS studies of AML genomes from children and adults have provided a comprehensive map of the genetic and epigenetic abnormalities, as well as the resulting transcriptional changes. Although, all the recurrent mutations in AML have at this time been identified, there are still some key questions needing to be answered, for instance, which mutations can initiate or contribute to the maintenance of AML? Which mutations cause drug resistance and which genes are responsible for relapse?

Therefore, the next big challenge in this field is to identify and functionally characterize genes and pathways with a causative role in the pathogenesis and progression of AML. This will provide better understanding of the underlying mechanisms of AML and thereby help in the development of novel therapeutic strategies. Functional screening including shRNA screening is a promising tool for effective identification of essential genes and pathways in AML. In this thesis, we confirmed the reliability of this technique to uncover crucial novel genes in leukemic cell growth. Moreover, we established a reliable model system and methods including *ex-vivo* expansion of AML and normal hematopoietic cells and intra-femoral transplantation of AML patient cells, to investigate the role of novel genes in AML disease and revealed the functional role of the selected candidates in leukemic cell growth. Hence, our methods might be helpful to shed light on these important questions.

Furthermore, another major reason for the high mortality rate is that few effective drugs have been developed to treat AML. The standard care of treatment, i.e. chemotherapy, has remained mainly unimproved over the past 40 years. During the last decade, numerous studies have been performed to develop new drugs for the treatment of AML but more than 90% of them have failed in the clinical phase. The major problem is that these strategies have not been efficient to completely eliminate AML patient cells and achieve CR or they cause severe side effects on normal cells. Taken together, in AML disease, novel treatment strategies are essentially needed to improve the prognosis for these patients.

One major reason why many new treatments against AML have been failed in clinical trials is due to the intra- and inter-tumor heterogeneity of AML. Moreover, LICs, responsible for the emergence of the disease and relapse after treatment, are very rare in AML cells derived from patients. These are factors that must be considered in any drug discovery research program to provide efficient treatments. In contrast, pharmaceutical companies are typically

limited to using cell lines with poor predictive capabilities, without testing on relevant normal cell based models in pre-clinical trials, which leads to high failure rates in phase I/II clinical trials due to low efficacy and high toxicity profiles of new pharmaceutical compounds.

Since drug development strategies are very time and cost consuming, it might be worth to screen and validate the promising compound candidates first on AML patient samples in particular LICs and normal hematopoietic cells in early stages of drug development then continue with more reliable candidates. Our validation *in vitro* and *in vivo* methods may be a good resource to select more effective targets and new drugs in AML. In conclusion, for future drug development strategies in addition to use advanced techniques, it is important to include patient samples as well as relevant normal cells or tissue models to discover new efficient treatments with curative potential.

## 6 ACKNOWLEDGEMENTS

The most precious part of my PhD was to meet and spend time with extraordinary people. I want to express my sincere appreciation to all and especially thank those who most helped me during my PhD journey. It is crystal clear it would not be possible to make it without you.

Firstly, I would like to express my gratitude to my main supervisor **Julian Walfridsson** for giving me the opportunity to do my PhD on exciting projects. I really appreciate your consistent support and kindness, and all you taught me. Thank you for teaching me to be an independent scientist with a critical mind and also for your great knowledge and valuable advice. Thank you!

**Hong Qian**, my co-supervisor, thank you for all your help, generous support, wonderful feedback and enormous knowledge. I am grateful to you for considering me as one of your own students and teaching me many techniques. I learned a lot from your valuable skills and enjoyed discussing science with you. You are a great scientist.

My co-supervisor, **Petter Höglund**, thank you for your advice and support. It was great to collaborate with you and I really learned from your scientific thinking.

**Behnam Sadeghi**, my mentor, there are no words that could express my admiration and appreciation for all your kind support and inspiring me when I was disappointed. Thanks a million for your trust when I met you the first time and for helping me come to Sweden to do my PhD. I consider you as my older brother and will never forget your help.

I wish to express my deepest gratitude to **Moustapha Hassan**; thank you for inviting me to Sweden and to Karolinska Institutet.

A special thanks to **Julian's group** at HERM. To **Gözde Türköz**, for your help and hard work. I could not have done experiments without your help. You are the best! Also, thanks for your presents to Adrina and taking care of her when I was working in the virus lab. My colleague **Aditya Harisankar**, for your contribution in the projects and great working environment and also the fantastic times that we played football. **Shabnam Kharazi**, for all your kind help in the projects, your contribution to make movies and your surprises, especially delicious tuna fish. Many thanks for taking care of Adrina and for ensuring that she did not make a big mess in the lab. To all the Master students who helped me in the projects, especially **Emma Wagner** for your assistance in the projects and in particular your generous and kind help in revising my thesis. **David Chang**, **Dora Boukoura** and **Sara**

**Shahsavarian** thanks for your great work, **Esmat Kamali** thank you for your help, your kind-heartedness and providing a fun environment in cell culture. I appreciate all the scientific input and discussions that I received from my group members.

I would like to say a big thank you to all the bone marrow **donors** in my studies, kind **midwives** at Karolinska Huddinge for collecting umbilical cord blood and our fantastic colleagues at **PKL4** and **PKL5**. This work could have not been done without your help.

I would like to thank all our collaborators, co-authors in the articles and people who helped me in my projects and my scientific development. Our collaborators in SciLife; **Mikael Altun** and **Johan Boström** for kindly providing shRNA and CRISPR/Cas9 vectors. **Nikolas Herold**, for the SAMHD1 project. **Aleksandra Krstic**, for her kind assistance in preparing RNA sequencing samples. To the wonderful people at **Sprint Bioscience**, in particular **Johan Lindström**, **Andreas Höglund**, **Jenny Viklund**, **Jessica Martinsson** and **Martin Andersson** who taught me all the processes to develop a new compound and generously shared your experiences. To the kind and supportive people in **BEA**; **Fredrik Fagerström-Billai**, **Patrick Muller**, **Tassos Damdimopoulos**, **David Brodin**, **Marika Rönholm** and **Susann Fält**, for your help to perform RNA-seq, ChIP-seq and bioinformatics analysis. I know I always stressed you to run and analyze my samples as fast as possible!

As we learned the important role of the BM niche in hematopoiesis, I would say my working environment also had an undeniable positive impact on my scientific and personal development. I would like to express my gratitude to the brilliant people at one of the best places at KI, **HERM**. I enjoyed doing my PhD at HERM and meeting fantastic people very much. Each one of you has been part of my progress during my unforgettable PhD journey. I would like to express my heartfelt appreciation to all of you for sharing your expertise, and knowledge, and moments of happiness and sadness. Thanks to all the colleagues that made HERM a friendly environment to work in particular:

**Eva Hellström-Lindberg**, for the fantastic effort to establish HERM with a nice and friendly environment, and for all your kind support and help during my studies. To **Johanna Ungerstedt**, for being the chairperson at my defense. **Sören Lehman**, for collaboration and allowing access to samples. **Sten Eirik Jacobsen**, for giving great scientific comments and teaching us to think critically. **Robert Månsson**, **Evren Alici**, **Yenan Bryceson**, **Petter Woll** and **Sidinh Luc** for your scientific discussions, great comments at SAPs and Friday meetings. I learned a lot from you. **Iyadh Douagi**, who taught me flow cytometry and sorting techniques. You were always kind and patient to answer my questions even when you were



busy. **Lindsay Davies**, for all your help, you are very kind and supportive. I never forget you kindly accepting to revise my thesis while you were on vacation.

**Monika Jansson**, thank you for all your generous and kind help. You are one of the kindest people I have ever met and I cannot imagine how we could work at HERM without your non-stop efforts. **Jelve Zendegani**, we are lucky to have such a nice person at HERM. You are always with kind smiles and positive energy. Thanks for your kindness, your help and advice in stressful situations. **Sri Sahlin**, for very kind help and your smile anytime I knocked on your door. You are great and supportive. **Annette Öster Fernström**, thanks for all your help. **Charlotte Gustafsson**, for your positive energy. **Sofia Bengtzen**, for your kind help in accessing the samples. **Anne-Sofie Johansson**, for all your help, fun and singing songs. **Mari Gilljam**, **Lili Andersson** and **Gunilla Walldin** for being kind and supportive.

I warmly remember my great friends who shared time with me and I am honored to have you as friends: **Nadir Kadri**, my best colleague, great scientist and a true friend that I can always count on. We had a great time together at HERM and in our meeting place. Thanks for all the invitations, laughs, advice and breaking news! **Stephan Meinke**, for your precious help to revise my thesis, all after works, fun watching and playing football, dinners and most importantly all your assistance and creative ideas to make movies. **Marios Dimitriou**, for generous sharing of your experience and knowledge and teaching me LTC-IC method. **Simona Conte**, for great times in cell culture and after work and for teaching many useful Italian key words! Without you the lab was completely quiet. **Ying Qu** for your help and kind advice. I will never forget your great help in the hard time of submitting articles. Special thanks to **Hani Abdulkadir ali**, **Deepika Nair** and **Hero Nikdin** (sare piaz) for all the fantastic moments, enjoyable times and gossips! Hope we can meet each other in future.

My other friends: **Monika Dolinska** and **Makoto Konko** for all the fun and pleasant times especially in the USA. **Pingnan Xiao** for your help and fun during our collaboration and in the office. **Thuy Luu Thanh**, for your kindness and the great time in the office. **Saeed Eshtad** for nice discussions about science and life. **Ece Somuncular** for delicious Turkish sweets. **Ayla De Paepe** and **Lucia Pena Perez** for kindly answering my questions about bioinformatics. **Kari Högstrand** for answering and helping with questions regarding vectors and GMM. **Hongya Han** and **Lamberto Torralba Raga** for spending time in the office at old HERM and all the jokes. **Lakshmi Sandhow** for all your help and lending me stuff any time I asked. **Matthias Voss** for scientific discussions in the big molecular lab and answering questions about WB, **Giovanna Perinetti Casoni** for the enjoyable time to make the last

movie “History of HERM”, **Desmond Chin**, **Allegra Matheson Lord** and **Edyta Wójtowicz** for great time and talks every Friday. **Tolga Sutu** and **Adil Doganay Duro**, I miss you a lot. Hope to see you again. **Srinivas Rao Nagubothu**, **Fawaz abomaray**, **Sigrun Stulz**, **Irene Gutierrez Perez**, **Donatella Galgano**, **Beatrice Zitti**, **Thaibault Boudierlique**, **Teresa Mortera-Blanco**, **Caroline Leijonhufvud**, **Filip Segerberg**, **Alamdar Hussain**, **Michael Chrobok**, **Huthayfa Mujahed**, **Sridharan Ganesan**, **Arnika Kathleen Wagner** and **Caroline Gavin** for all the fun, helps, fika and discussions. Thank you all.

To the **Department of Medicine** especially: **Jan Bolinder**, **Klas Karlsson**, **Ulrika Markne**, **Anastasia Urban**, **Elenor Nyman**, **Therese Lind** and **Edgardo Faune** for all your help.

All members of the Moustapha Hassan group: **Ebrahim**, **Sulaiman**, **Mona** and **Maria**.  
Thanks for your time and teaching me to start to work in KI.

I would like to acknowledge those who have not been directly involved in my thesis but their presence and support helped me to enjoy and successfully achieve my PhD. My deepest heartfelt thanks to my **Iranian friends** in Sweden. I am lucky to have you as fantastic friends here, in particular:

**Bitā Khoein**, we had a difficult time during the first year in Sweden and when we didn't have a place to stay for a few days, you warmly invited us to stay with you. Anytime that we felt homesick, you always made time for us. I will never forget your kindness. Dear **Sahba**, our extremely kind friend, for making very nice birthday cards and my friend **Illia** for joining me to play football. **Ghazaleh Assadi**, from the first week that I came to Sweden until now, you have always helped me with everything I needed. I was lucky to meet you. Thanks for everything, for your wonderful and kind friendship, for all your support. Your heart is bigger than the sea. Also, I would like to thank your family especially your parents. **Maryam Nikpour**, I remember the first day that I arrived in Stockholm and I was not even confident to do shopping. You showed me how to shop and we had the first tour at Novum. **Roger Hult**, **Sofia**, **Sam** and little **Simon** thanks a lot for all your kindness during these years. **Mehdi** and **Nasrin Siyahtiri** I have no words to express how grateful I am for being my family in Sweden. I consider you as my brother and sister. You were always there in the most difficult times and helped me regardless of when I needed you. **Mastoureh Shahsavani** and **Shahin Akbarnezhad**, we had great moments together especially our camping. We spend most of the time with you, we celebrate many events together and we always enjoyed to meet you. Thank you for being the best friends, your warm hospitality and perfect cooking from Mastoureh Banoo! Thanks to little princess **Diana** for being one of the best friends to Adrina.

**Nasim Derakhshan** and **Amir Moulavi**, for being wonderful, supportive and true friends and for all the fun moments that we had together. Dear **Borna**, thanks for your generosity and kindness sharing toys with Adrina. **Mohsen Karimi**, **Atosa**, **Parnian** and **Ardalan**, you are great friends. We had unforgettable moments celebrating Persian events together and I will never forget our memorable trip to Denmark. Mohsen thanks for all your kind help, grilling and your lovely Kebabs. **Manuchehr Abedi**, always calm, handsome and a gentleman. Thanks for your generous advice. **Masoud Vosough**, thank you for being such a nice friend and I hope to meet you more often. **Mona Noohi**, my nice and kind friend. I would never forget your Lobia polo and I found you as a potential candidate to babysit Adrina! **Laia Sadeghi** and **Siamak Akbari** for your warm hospitality and great moments that we spend together. **Zahra Golabkesh** and **Behrouz Afsharian**, for being nice friends and sharing time with us. **Maryam Saghafian** and **Raul**, I am grateful to know you. Thanks for great homemade pasta, all the scientific discussions and outdoor activities. **Yasaman Zahedi**, for your kindness and always being supportive. I will never forget your delicious chocolate cakes and fantastic Ghalye Mahi. **Farzaneh Shahin** for being a good friend, your delicious surprises and your kindness that I will never forget. **Marjan Abbasi** for the brilliant times together and all big laughs with apple taste. **Amir Mahani** for the jokes and enjoyable times watching football matches together. **Maryam Iravani** and **Konstantin**, for your delicious Shole zard and Fika at your place. **Mahyar** and **Mohammad** for being good friends. **Marzieh Saeedimasine** for being new neighbor and nice friend in NEO.

All my **teachers** and **supervisors** who helped me during my studies in Iran, in particular the symbol of morality Professor **Moulavi**.

My dear friends in Iran: my dear cousin **Mehdi Heshmati**, I would never feel satisfied even if I visited you every day. Thanks for making all the unforgettable moments with me. **Behrouz Salamt** and **Parivash Khodayar**, my dear friends. I have always enjoyed your company. My best friend **Javad Ansari**, anytime that I called you, you accepted to come and visit me. Thank you so much. **Mahmoud Haghverdi**, **Farshid Alaedini** for being a kind friends and spending time together. My previous colleagues in **Welfare and Rehabilitation center** and **Forensic Genetics**, thanks for being supportive and kind friends.

I would like to thank the most important people in my life; **my family**, my parents **Maman** and **Baba**:

پدر و مادر عزیزم، تشکر را بپذیرید بابت تمام زحمات و سختی‌هایی که در طول سال‌ها تحمل کردید تا ما سرانجام به موفقیت دست یابیم. هرگز تلاش و محبت شما را از خاطر نخواهم برد. دوستتان دارم.

My great **mother** and **father in law**, many thanks for your continuous support and kindness. Also thanks my brothers in law **Farhad** and **Jafar** and my sisters in law **Marjan**, **Shahin** and **Roghayeh**. My brother **Naser** and my sister **Mahsa**:

برادر عزیزم ناصر و خواهر مهربانم مهسا، سپاس برای تمام مهربانی‌ها و خاطرات زیبا. همیشه سلامت شاد و خوشبخت باشید.

My sisters **Sara** and **Zahra**:

خواهران عزیزم، بهترین‌ها و زیباترین لحظه‌های زندگی را برای شما آرزومندم.

My dear love and best friend **Manizheh**, I would like to thank you the most, as the most important person who helped me during my PhD, for all your support and help during these years. It is very clear that without your help I could not have achieved my PhD. We met each other 16 years back in Ahvaz when we were Bachelor students. We decided to proceed in science and make a future together. During these years, we had hard times and great moments. Thanks for your patience during many weekends and nights that I was busy in the lab and you were alone at home. I love you so much from bottom of my heart!

My little angel, **Adrina**, I don't know when you can read this by yourself. You don't know how much the world became brighter, meaningful and wonderful when you came into my life. Thank you for accepting me to be your father. You are my "Nafas" and I will love you forever.

*Life is the unique scene of our artistry  
Everyone sings his own song and leaves the scene  
The scene is constant  
Blessed the song that people bear it in mind  
Zhaleh Esfehani (1921-2007)*

This thesis was supported by Wallenberg Foundation, Åke Wiberg Foundation, Åke Olsson Foundation for Hematological research, Magnus Bergvall Foundation, Karolinska Institutet, Swedish Cancer Society, Vinnova, The Swedish Childhood Cancer Foundation.

## 7 REFERENCES

- Abdel-Wahab O, Adli M, LaFave LM, Gao J, Hricik T, Shih AH, et al. ASXL1 mutations promote myeloid transformation through loss of PRC2-mediated gene repression. *Cancer Cell*. 2012;22(2):180-93.
- Adan A, Alizada G, Kiraz Y, Baran Y, Nalbant A. Flow cytometry: basic principles and applications. *Crit Rev Biotechnol*. 2017;37(2):163-76.
- Adli M. The CRISPR tool kit for genome editing and beyond. *Nat Commun*. 2018;9(1):1911.
- Adolfsson J, Mansson R, Buza-Vidas N, Hultquist A, Liuba K, Jensen CT, et al. Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell*. 2005;121(2):295-306.
- Akasaka T, Tsuji K, Kawahira H, Kanno M, Harigaya K, Hu L, et al. The role of mel-18, a mammalian Polycomb group gene, during IL-7-dependent proliferation of lymphocyte precursors. *Immunity*. 1997;7(1):135-46.
- Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SA, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. *Nature*. 2013;500(7463):415-21.
- Alharbi RA, Pettengell R, Pandha HS, Morgan R. The role of HOX genes in normal hematopoiesis and acute leukemia. *Leukemia*. 2013;27(5):1000-8.
- Altman R, Kellogg D. Control of mitotic events by Nap1 and the Gin4 kinase. *J Cell Biol*. 1997;138(1):119-30.
- Amadori S, Suciu S, Selleslag D, Aversa F, Gaidano G, Musso M, et al. Gemtuzumab Ozogamicin Versus Best Supportive Care in Older Patients With Newly Diagnosed Acute Myeloid Leukemia Unsuitable for Intensive Chemotherapy: Results of the Randomized Phase III EORTC-GIMEMA AML-19 Trial. *J Clin Oncol*. 2016;34(9):972-9.
- Appelbaum FR, Bernstein ID. Gemtuzumab ozogamicin for acute myeloid leukemia. *Blood*. 2017;130(22):2373-6.
- Arber DA, Orazi A, Hasserjian R, Thiele J, Borowitz MJ, Le Beau MM, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-405.
- Aubrey BJ, Kelly GL, Kueh AJ, Brennan MS, O'Connor L, Milla L, et al. An inducible lentiviral guide RNA platform enables the identification of tumor-essential genes and tumor-promoting mutations in vivo. *Cell Rep*. 2015;10(8):1422-32.
- Bagger FO, Sasivarevic D, Sohi SH, Laursen LG, Pundhir S, Sonderby CK, et al. BloodSpot: a database of gene expression profiles and transcriptional programs for healthy and malignant haematopoiesis. *Nucleic Acids Res*. 2016;44(D1):D917-24.
- Bar-Natan M, Nelson EA, Xiang M, Frank DA. STAT signaling in the pathogenesis and treatment of myeloid malignancies. *JAKSTAT*. 2012;1(2):55-64.
- Baum CM, Weissman IL, Tsukamoto AS, Buckle AM, Peault B. Isolation of a candidate human hematopoietic stem-cell population. *Proc Natl Acad Sci U S A*. 1992;89(7):2804-8.

- Beerman I, Maloney WJ, Weissmann IL, Rossi DJ. Stem cells and the aging hematopoietic system. *Curr Opin Immunol.* 2010;22(4):500-6.
- Belaoussoff M, Farrington SM, Baron MH. Hematopoietic induction and respecification of A-P identity by visceral endoderm signaling in the mouse embryo. *Development.* 1998;125(24):5009-18.
- Bennett JM, Catovsky D, Daniel MT, Flandrin G, Galton DA, Gralnick HR, et al. Proposals for the classification of the acute leukaemias. French-American-British (FAB) co-operative group. *Br J Haematol.* 1976;33(4):451-8.
- Bhatia M, Wang JC, Kapp U, Bonnet D, Dick JE. Purification of primitive human hematopoietic cells capable of repopulating immune-deficient mice. *Proc Natl Acad Sci U S A.* 1997;94(10):5320-5.
- Boissel N, Leroy H, Brethon B, Philippe N, de Botton S, Auvrignon A, et al. Incidence and prognostic impact of c-Kit, FLT3, and Ras gene mutations in core binding factor acute myeloid leukemia (CBF-AML). *Leukemia.* 2006;20(6):965-70.
- Bowman RL, Busque L, Levine RL. Clonal Hematopoiesis and Evolution to Hematopoietic Malignancies. *Cell Stem Cell.* 2018;22(2):157-70.
- Brooke-Bisschop T, Savory JGA, Foley T, Ringuette R, Lohnes D. Essential roles for Cdx in murine primitive hematopoiesis. *Dev Biol.* 2017;422(2):115-24.
- Bullinger L, Dohner K, Dohner H. Genomics of Acute Myeloid Leukemia Diagnosis and Pathways. *J Clin Oncol.* 2017;35(9):934-46.
- Burnett AK, Russell NH, Hunter AE, Milligan D, Knapper S, Wheatley K, et al. Clofarabine doubles the response rate in older patients with acute myeloid leukemia but does not improve survival. *Blood.* 2013;122(8):1384-94.
- Burns CE, Traver D, Mayhall E, Shepard JL, Zon LI. Hematopoietic stem cell fate is established by the Notch-Runx pathway. *Genes Dev.* 2005;19(19):2331-42.
- Busque L, Patel JP, Figueroa ME, Vasanthakumar A, Provost S, Hamilou Z, et al. Recurrent somatic TET2 mutations in normal elderly individuals with clonal hematopoiesis. *Nat Genet.* 2012;44(11):1179-81.
- Byrd JC, Mrozek K, Dodge RK, Carroll AJ, Edwards CG, Arthur DC, et al. Pretreatment cytogenetic abnormalities are predictive of induction success, cumulative incidence of relapse, and overall survival in adult patients with de novo acute myeloid leukemia: results from Cancer and Leukemia Group B (CALGB 8461). *Blood.* 2002;100(13):4325-36.
- Caligiuri MA, Schichman SA, Strout MP, Mrozek K, Baer MR, Frankel SR, et al. Molecular rearrangement of the ALL-1 gene in acute myeloid leukemia without cytogenetic evidence of 11q23 chromosomal translocations. *Cancer Res.* 1994;54(2):370-3.
- Caligiuri MA, Strout MP, Lawrence D, Arthur DC, Baer MR, Yu F, et al. Rearrangement of ALL1 (MLL) in acute myeloid leukemia with normal cytogenetics. *Cancer Res.* 1998;58(1):55-9.
- Calvi LM, Adams GB, Weibrecht KW, Weber JM, Olson DP, Knight MC, et al. Osteoblastic cells regulate the haematopoietic stem cell niche. *Nature.* 2003;425(6960):841-6.
- Campos L, Guyotat D, Archimbaud E, Devaux Y, Treille D, Larese A, et al. Surface marker expression in adult acute myeloid leukaemia: correlations with initial characteristics, morphology and response to therapy. *Br J Haematol.* 1989;72(2):161-6.

- Cancer Genome Atlas Research N, Ley TJ, Miller C, Ding L, Raphael BJ, Mungall AJ, et al. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med*. 2013;368(22):2059-74.
- Carrelha J, Meng Y, Kettle LM, Luis TC, Norfo R, Alcolea V, et al. Hierarchically related lineage-restricted fates of multipotent haematopoietic stem cells. *Nature*. 2018;554(7690):106-11.
- Casero D, Sandoval S, Seet CS, Scholes J, Zhu Y, Ha VL, et al. Long non-coding RNA profiling of human lymphoid progenitor cells reveals transcriptional divergence of B cell and T cell lineages. *Nat Immunol*. 2015;16(12):1282-91.
- Catlin SN, Busque L, Gale RE, Gutter P, Abkowitz JL. The replication rate of human hematopoietic stem cells in vivo. *Blood*. 2011;117(17):4460-6.
- Challen GA, Sun D, Mayle A, Jeong M, Luo M, Rodriguez B, et al. Dnmt3a and Dnmt3b have overlapping and distinct functions in hematopoietic stem cells. *Cell Stem Cell*. 2014;15(3):350-64.
- Chen CZ, Li L, Lodish HF, Bartel DP. MicroRNAs modulate hematopoietic lineage differentiation. *Science*. 2004;303(5654):83-6.
- Cheshier SH, Morrison SJ, Liao X, Weissman IL. In vivo proliferation and cell cycle kinetics of long-term self-renewing hematopoietic stem cells. *Proc Natl Acad Sci U S A*. 1999;96(6):3120-5.
- Cheson BD, Bennett JM, Kopecky KJ, Buchner T, Willman CL, Estey EH, et al. Revised recommendations of the International Working Group for Diagnosis, Standardization of Response Criteria, Treatment Outcomes, and Reporting Standards for Therapeutic Trials in Acute Myeloid Leukemia. *J Clin Oncol*. 2003;21(24):4642-9.
- Chou WC, Chou SC, Liu CY, Chen CY, Hou HA, Kuo YY, et al. TET2 mutation is an unfavorable prognostic factor in acute myeloid leukemia patients with intermediate-risk cytogenetics. *Blood*. 2011;118(14):3803-10.
- Chudnovsky Y, Kim D, Zheng S, Whyte WA, Bansal M, Bray MA, et al. ZFH4 interacts with the NuRD core member CHD4 and regulates the glioblastoma tumor-initiating cell state. *Cell Rep*. 2014;6(2):313-24.
- Cimmino L, Dolgalev I, Wang Y, Yoshimi A, Martin GH, Wang J, et al. Restoration of TET2 Function Blocks Aberrant Self-Renewal and Leukemia Progression. *Cell*. 2017;170(6):1079-95 e20.
- Civin CI, Strauss LC, Brovall C, Fackler MJ, Schwartz JF, Shaper JH. Antigenic analysis of hematopoiesis. III. A hematopoietic progenitor cell surface antigen defined by a monoclonal antibody raised against KG-1a cells. *J Immunol*. 1984;133(1):157-65.
- Clifford R, Louis T, Robbe P, Ackroyd S, Burns A, Timbs AT, et al. SAMHD1 is mutated recurrently in chronic lymphocytic leukemia and is involved in response to DNA damage. *Blood*. 2014;123(7):1021-31.
- Cooper TM, Franklin J, Gerbing RB, Alonzo TA, Hurwitz C, Raimondi SC, et al. AAML03P1, a pilot study of the safety of gemtuzumab ozogamicin in combination with chemotherapy for newly diagnosed childhood acute myeloid leukemia: a report from the Children's Oncology Group. *Cancer*. 2012;118(3):761-9.

- Corces-Zimmerman MR, Hong WJ, Weissman IL, Medeiros BC, Majeti R. Preleukemic mutations in human acute myeloid leukemia affect epigenetic regulators and persist in remission. *Proc Natl Acad Sci U S A*. 2014;111(7):2548-53.
- Cozzio A, Passegue E, Ayton PM, Karsunky H, Cleary ML, Weissman IL. Similar MLL-associated leukemias arising from self-renewing stem cells and short-lived myeloid progenitors. *Genes Dev*. 2003;17(24):3029-35.
- Crispino JD, Horwitz MS. GATA factor mutations in hematologic disease. *Blood*. 2017;129(15):2103-10.
- Cui Y, Xu J, Cheng M, Liao X, Peng S. Review of CRISPR/Cas9 sgRNA Design Tools. *Interdiscip Sci*. 2018;10(2):455-65.
- Dahl R, Iyer SR, Owens KS, Cuylear DD, Simon MC. The transcriptional repressor GFI-1 antagonizes PU.1 activity through protein-protein interaction. *J Biol Chem*. 2007;282(9):6473-83.
- De Kouchkovsky I, Abdul-Hay M. 'Acute myeloid leukemia: a comprehensive review and 2016 update'. *Blood Cancer J*. 2016;6(7):e441.
- de Pater E, Kaimakis P, Vink CS, Yokomizo T, Yamada-Inagawa T, van der Linden R, et al. Gata2 is required for HSC generation and survival. *J Exp Med*. 2013;210(13):2843-50.
- Deveau H, Garneau JE, Moineau S. CRISPR/Cas system and its role in phage-bacteria interactions. *Annu Rev Microbiol*. 2010;64:475-93.
- DiNardo CD, Bannon SA, Routbort M, Franklin A, Mork M, Armanios M, et al. Evaluation of Patients and Families With Concern for Predispositions to Hematologic Malignancies Within the Hereditary Hematologic Malignancy Clinic (HHMC). *Clin Lymphoma Myeloma Leuk*. 2016;16(7):417-28 e2.
- Ding L, Ley TJ, Larson DE, Miller CA, Koboldt DC, Welch JS, et al. Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature*. 2012;481(7382):506-10.
- Dohner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, Buchner T, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*. 2017;129(4):424-47.
- Dohner H, Estey EH, Amadori S, Appelbaum FR, Buchner T, Burnett AK, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*. 2010;115(3):453-74.
- Dohner H, Weisdorf DJ, Bloomfield CD. Acute Myeloid Leukemia. *N Engl J Med*. 2015;373(12):1136-52.
- Dohner K, Schlenk RF, Habdank M, Scholl C, Rucker FG, Corbacioglu A, et al. Mutant nucleophosmin (NPM1) predicts favorable prognosis in younger adults with acute myeloid leukemia and normal cytogenetics: interaction with other gene mutations. *Blood*. 2005;106(12):3740-6.
- Dores GM, Devesa SS, Curtis RE, Linet MS, Morton LM. Acute leukemia incidence and patient survival among children and adults in the United States, 2001-2007. *Blood*. 2012;119(1):34-43.



- Doulatov S, Notta F, Laurenti E, Dick JE. Hematopoiesis: a human perspective. *Cell Stem Cell*. 2012;10(2):120-36.
- Dykstra B, Kent D, Bowie M, McCaffrey L, Hamilton M, Lyons K, et al. Long-term propagation of distinct hematopoietic differentiation programs in vivo. *Cell Stem Cell*. 2007;1(2):218-29.
- Dzierzak E, Bigas A. Blood Development: Hematopoietic Stem Cell Dependence and Independence. *Cell Stem Cell*. 2018;22(5):639-51.
- Dzierzak E, Medvinsky A. Mouse embryonic hematopoiesis. *Trends Genet*. 1995;11(9):359-66.
- Ebralidze AK, Guibal FC, Steidl U, Zhang P, Lee S, Bartholdy B, et al. PU.1 expression is modulated by the balance of functional sense and antisense RNAs regulated by a shared cis-regulatory element. *Genes Dev*. 2008;22(15):2085-92.
- Eppert K, Takenaka K, Lechman ER, Waldron L, Nilsson B, van Galen P, et al. Stem cell gene expression programs influence clinical outcome in human leukemia. *Nat Med*. 2011;17(9):1086-93.
- Estey E, Dohner H. Acute myeloid leukaemia. *Lancet*. 2006;368(9550):1894-907.
- Estey EH. Acute myeloid leukemia: 2014 update on risk-stratification and management. *Am J Hematol*. 2014;89(11):1063-81.
- Etchin J, Sun Q, Kentsis A, Farmer A, Zhang ZC, Sanda T, et al. Antileukemic activity of nuclear export inhibitors that spare normal hematopoietic cells. *Leukemia*. 2013;27(1):66-74.
- Fabian MR, Sonenberg N. The mechanics of miRNA-mediated gene silencing: a look under the hood of miRISC. *Nat Struct Mol Biol*. 2012;19(6):586-93.
- Faderl S, Ravandi F, Huang X, Garcia-Manero G, Ferrajoli A, Estrov Z, et al. A randomized study of clofarabine versus clofarabine plus low-dose cytarabine as front-line therapy for patients aged 60 years and older with acute myeloid leukemia and high-risk myelodysplastic syndrome. *Blood*. 2008;112(5):1638-45.
- Falini B, Nicoletti I, Martelli MF, Mecucci C. Acute myeloid leukemia carrying cytoplasmic/mutated nucleophosmin (NPMc+ AML): biologic and clinical features. *Blood*. 2007;109(3):874-85.
- Fasan A, Haferlach C, Alpermann T, Jeromin S, Grossmann V, Eder C, et al. The role of different genetic subtypes of CEBPA mutated AML. *Leukemia*. 2014;28(4):794-803.
- Fathi AT, Wander SA, Faramand R, Emadi A. Biochemical, Epigenetic, and Metabolic Approaches to Target IDH Mutations in Acute Myeloid Leukemia. *Semin Hematol*. 2015;52(3):165-71.
- Fernandez HF, Sun Z, Yao X, Litzow MR, Luger SM, Paietta EM, et al. Anthracycline dose intensification in acute myeloid leukemia. *N Engl J Med*. 2009;361(13):1249-59.
- Ferreira HJ, Heyn H, Vizoso M, Moutinho C, Vidal E, Gomez A, et al. DNMT3A mutations mediate the epigenetic reactivation of the leukemogenic factor MEIS1 in acute myeloid leukemia. *Oncogene*. 2016;35(23):3079-82.
- Ferrell JE, Jr. Bistability, bifurcations, and Waddington's epigenetic landscape. *Curr Biol*. 2012;22(11):R458-66.

- Figuerola ME, Abdel-Wahab O, Lu C, Ward PS, Patel J, Shih A, et al. Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell*. 2010;18(6):553-67.
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*. 1998;391(6669):806-11.
- Fischbach NA, Rozenfeld S, Shen W, Fong S, Chrobak D, Ginzinger D, et al. HOXB6 overexpression in murine bone marrow immortalizes a myelomonocytic precursor in vitro and causes hematopoietic stem cell expansion and acute myeloid leukemia in vivo. *Blood*. 2005;105(4):1456-66.
- Fujiwara Y, Browne CP, Cunniff K, Goff SC, Orkin SH. Arrested development of embryonic red cell precursors in mouse embryos lacking transcription factor GATA-1. *Proc Natl Acad Sci U S A*. 1996;93(22):12355-8.
- Fukuda M, Asano S, Nakamura T, Adachi M, Yoshida M, Yanagida M, et al. CRM1 is responsible for intracellular transport mediated by the nuclear export signal. *Nature*. 1997;390(6657):308-11.
- Gaidzik VI, Paschka P, Spath D, Habdank M, Kohne CH, Germing U, et al. TET2 mutations in acute myeloid leukemia (AML): results from a comprehensive genetic and clinical analysis of the AML study group. *J Clin Oncol*. 2012;30(12):1350-7.
- Gaidzik VI, Schlenk RF, Paschka P, Stolzle A, Spath D, Kuendgen A, et al. Clinical impact of DNMT3A mutations in younger adult patients with acute myeloid leukemia: results of the AML Study Group (AMLSG). *Blood*. 2013;121(23):4769-77.
- Gale AJ. Continuing education course #2: current understanding of hemostasis. *Toxicol Pathol*. 2011;39(1):273-80.
- Gale RE, Green C, Allen C, Mead AJ, Burnett AK, Hills RK, et al. The impact of FLT3 internal tandem duplication mutant level, number, size, and interaction with NPM1 mutations in a large cohort of young adult patients with acute myeloid leukemia. *Blood*. 2008;111(5):2776-84.
- Galloway JL, Zon LI. Ontogeny of hematopoiesis: examining the emergence of hematopoietic cells in the vertebrate embryo. *Curr Top Dev Biol*. 2003;53:139-58.
- Gamis AS, Alonzo TA, Perentesis JP, Meshinchi S, Committee COGAML. Children's Oncology Group's 2013 blueprint for research: acute myeloid leukemia. *Pediatr Blood Cancer*. 2013;60(6):964-71.
- Gao S, Yang C, Jiang S, Xu XN, Lu X, He YW, et al. Applications of RNA interference high-throughput screening technology in cancer biology and virology. *Protein Cell*. 2014;5(11):805-15.
- Geisler S, Collier J. RNA in unexpected places: long non-coding RNA functions in diverse cellular contexts. *Nat Rev Mol Cell Biol*. 2013;14(11):699-712.
- Genovese G, Kahler AK, Handsaker RE, Lindberg J, Rose SA, Bakhoum SF, et al. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med*. 2014;371(26):2477-87.
- Gill S, Tasian SK, Ruella M, Shestova O, Li Y, Porter DL, et al. Preclinical targeting of human acute myeloid leukemia and myeloablation using chimeric antigen receptor-modified T cells. *Blood*. 2014;123(15):2343-54.

- Gilliland DG, Griffin JD. The roles of FLT3 in hematopoiesis and leukemia. *Blood*. 2002;100(5):1532-42.
- Goardon N, Marchi E, Atzberger A, Quek L, Schuh A, Soneji S, et al. Coexistence of LMPP-like and GMP-like leukemia stem cells in acute myeloid leukemia. *Cancer Cell*. 2011;19(1):138-52.
- Graf T. Differentiation plasticity of hematopoietic cells. *Blood*. 2002;99(9):3089-101.
- Grande M, Lambea E, Fajardo A, Lopez-Aviles S, Kellogg D, Aligue R. Crosstalk between Nap1 protein and Cds1 checkpoint kinase to maintain chromatin integrity. *Biochim Biophys Acta*. 2008;1783(9):1595-604.
- Grass JA, Boyer ME, Pal S, Wu J, Weiss MJ, Bresnick EH. GATA-1-dependent transcriptional repression of GATA-2 via disruption of positive autoregulation and domain-wide chromatin remodeling. *Proc Natl Acad Sci U S A*. 2003;100(15):8811-6.
- Griessinger E, Anjos-Afonso F, Pizzitola I, Rouault-Pierre K, Vargaftig J, Taussig D, et al. A niche-like culture system allowing the maintenance of primary human acute myeloid leukemia-initiating cells: a new tool to decipher their chemoresistance and self-renewal mechanisms. *Stem Cells Transl Med*. 2014;3(4):520-9.
- Grimwade D, Freeman SD. Defining minimal residual disease in acute myeloid leukemia: which platforms are ready for "prime time"? *Blood*. 2014;124(23):3345-55.
- Grimwade D, Ivey A, Huntly BJ. Molecular landscape of acute myeloid leukemia in younger adults and its clinical relevance. *Blood*. 2016;127(1):29-41.
- Grove CS, Vassiliou GS. Acute myeloid leukaemia: a paradigm for the clonal evolution of cancer? *Dis Model Mech*. 2014;7(8):941-51.
- Guiu J, Shimizu R, D'Altri T, Fraser ST, Hatakeyama J, Bresnick EH, et al. Hes repressors are essential regulators of hematopoietic stem cell development downstream of Notch signaling. *J Exp Med*. 2013;210(1):71-84.
- Gurtner GC, Werner S, Barrandon Y, Longaker MT. Wound repair and regeneration. *Nature*. 2008;453(7193):314-21.
- Haferlach C, Dicker F, Herholz H, Schnittger S, Kern W, Haferlach T. Mutations of the TP53 gene in acute myeloid leukemia are strongly associated with a complex aberrant karyotype. *Leukemia*. 2008;22(8):1539-41.
- Hansen KH, Bracken AP, Pasini D, Dietrich N, Gehani SS, Monrad A, et al. A model for transmission of the H3K27me3 epigenetic mark. *Nat Cell Biol*. 2008;10(11):1291-300.
- Harrison DE. Mouse erythropoietic stem cell lines function normally 100 months: loss related to number of transplantations. *Mech Ageing Dev*. 1979;9(5-6):427-33.
- Heath EM, Chan SM, Minden MD, Murphy T, Shlush LI, Schimmer AD. Biological and clinical consequences of NPM1 mutations in AML. *Leukemia*. 2017;31(4):798-807.
- Hills RK, Castaigne S, Appelbaum FR, Delaunay J, Petersdorf S, Othus M, et al. Addition of gemtuzumab ozogamicin to induction chemotherapy in adult patients with acute myeloid leukaemia: a meta-analysis of individual patient data from randomised controlled trials. *Lancet Oncol*. 2014;15(9):986-96.

- Ho PA, Alonzo TA, Kopecky KJ, Miller KL, Kuhn J, Zeng R, et al. Molecular alterations of the IDH1 gene in AML: a Children's Oncology Group and Southwest Oncology Group study. *Leukemia*. 2010;24(5):909-13.
- Ho PA, Kutny MA, Alonzo TA, Gerbing RB, Joaquin J, Raimondi SC, et al. Leukemic mutations in the methylation-associated genes DNMT3A and IDH2 are rare events in pediatric AML: a report from the Children's Oncology Group. *Pediatr Blood Cancer*. 2011;57(2):204-9.
- Ho-Tin-Noe B. The multifaceted roles of platelets in inflammation and innate immunity. *Platelets*. 2018:1-2.
- Holliday R, Pugh JE. DNA modification mechanisms and gene activity during development. *Science*. 1975;187(4173):226-32.
- Huntly BJ, Shigematsu H, Deguchi K, Lee BH, Mizuno S, Duclos N, et al. MOZ-TIF2, but not BCR-ABL, confers properties of leukemic stem cells to committed murine hematopoietic progenitors. *Cancer Cell*. 2004;6(6):587-96.
- Ichikawa M, Asai T, Saito T, Seo S, Yamazaki I, Yamagata T, et al. AML-1 is required for megakaryocytic maturation and lymphocytic differentiation, but not for maintenance of hematopoietic stem cells in adult hematopoiesis. *Nat Med*. 2004;10(3):299-304.
- Ichikawa M, Yoshimi A, Nakagawa M, Nishimoto N, Watanabe-Okochi N, Kurokawa M. A role for RUNX1 in hematopoiesis and myeloid leukemia. *Int J Hematol*. 2013;97(6):726-34.
- Ikuta K, Weissman IL. Evidence that hematopoietic stem cells express mouse c-kit but do not depend on steel factor for their generation. *Proc Natl Acad Sci U S A*. 1992;89(4):1502-6.
- Ishii M, Matsuoka Y, Sasaki Y, Nakatsuka R, Takahashi M, Nakamoto T, et al. Development of a high-resolution purification method for precise functional characterization of primitive human cord blood-derived CD34-negative SCID-repopulating cells. *Exp Hematol*. 2011;39(2):203-13 e1.
- Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014;371(26):2488-98.
- Ji H, Ehrlich LI, Seita J, Murakami P, Doi A, Lindau P, et al. Comprehensive methylome map of lineage commitment from haematopoietic progenitors. *Nature*. 2010;467(7313):338-42.
- Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*. 2012;337(6096):816-21.
- Kajiume T, Ninomiya Y, Ishihara H, Kanno R, Kanno M. Polycomb group gene mel-18 modulates the self-renewal activity and cell cycle status of hematopoietic stem cells. *Exp Hematol*. 2004;32(6):571-8.
- Kantarjian HM, Erba HP, Claxton D, Arellano M, Lyons RM, Kovascovics T, et al. Phase II study of clofarabine monotherapy in previously untreated older adults with acute myeloid leukemia and unfavorable prognostic factors. *J Clin Oncol*. 2010;28(4):549-55.

- Kavanagh S, Murphy T, Law A, Yehudai D, Ho JM, Chan S, et al. Emerging therapies for acute myeloid leukemia: translating biology into the clinic. *JCI Insight*. 2017;2(18).
- Kayser S, Schlenk RF, Londono MC, Breitenbuecher F, Wittke K, Du J, et al. Insertion of FLT3 internal tandem duplication in the tyrosine kinase domain-1 is associated with resistance to chemotherapy and inferior outcome. *Blood*. 2009;114(12):2386-92.
- Kelly LM, Liu Q, Kutok JL, Williams IR, Boulton CL, Gilliland DG. FLT3 internal tandem duplication mutations associated with human acute myeloid leukemias induce myeloproliferative disease in a murine bone marrow transplant model. *Blood*. 2002;99(1):310-8.
- Kenderian SS, Ruella M, Shestova O, Klichinsky M, Aikawa V, Morrisette JJ, et al. CD33-specific chimeric antigen receptor T cells exhibit potent preclinical activity against human acute myeloid leukemia. *Leukemia*. 2015;29(8):1637-47.
- Kiel MJ, Morrison SJ. Maintaining hematopoietic stem cells in the vascular niche. *Immunity*. 2006;25(6):862-4.
- Kiel MJ, Yilmaz OH, Iwashita T, Yilmaz OH, Terhorst C, Morrison SJ. SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell*. 2005;121(7):1109-21.
- Kihara R, Nagata Y, Kiyoi H, Kato T, Yamamoto E, Suzuki K, et al. Comprehensive analysis of genetic alterations and their prognostic impacts in adult acute myeloid leukemia patients. *Leukemia*. 2014;28(8):1586-95.
- Kim I, Saunders TL, Morrison SJ. Sox17 dependence distinguishes the transcriptional regulation of fetal from adult hematopoietic stem cells. *Cell*. 2007;130(3):470-83.
- Kim SI, Bresnick EH. Transcriptional control of erythropoiesis: emerging mechanisms and principles. *Oncogene*. 2007;26(47):6777-94.
- Kindler T, Breitenbuecher F, Marx A, Beck J, Hess G, Weinkauff B, et al. Efficacy and safety of imatinib in adult patients with c-kit-positive acute myeloid leukemia. *Blood*. 2004;103(10):3644-54.
- Kingston RE, Chen CA, Rose JK. Calcium phosphate transfection. *Curr Protoc Mol Biol*. 2003;Chapter 9:Unit 9 1.
- Kobayashi H, Butler JM, O'Donnell R, Kobayashi M, Ding BS, Bonner B, et al. Angiocrine factors from Akt-activated endothelial cells balance self-renewal and differentiation of haematopoietic stem cells. *Nat Cell Biol*. 2010;12(11):1046-56.
- Kode A, Manavalan JS, Mosialou I, Bhagat G, Rathinam CV, Luo N, et al. Leukaemogenesis induced by an activating beta-catenin mutation in osteoblasts. *Nature*. 2014;506(7487):240-4.
- Kojima K, Kornblau SM, Ruvolo V, Dilip A, Duvvuri S, Davis RE, et al. Prognostic impact and targeting of CRM1 in acute myeloid leukemia. *Blood*. 2013;121(20):4166-74.
- Kosan C, Godmann M. Genetic and Epigenetic Mechanisms That Maintain Hematopoietic Stem Cell Function. *Stem Cells Int*. 2016;2016:5178965.
- Koschmieder S, Halmos B, Levantini E, Tenen DG. Dysregulation of the C/EBPalpha differentiation pathway in human cancer. *J Clin Oncol*. 2009;27(4):619-28.
- Kouzarides T. Chromatin modifications and their function. *Cell*. 2007;128(4):693-705.

- Krause DS, Fulzele K, Catic A, Sun CC, Dombkowski D, Hurley MP, et al. Differential regulation of myeloid leukemias by the bone marrow microenvironment. *Nat Med*. 2013;19(11):1513-7.
- Kreso A, Dick JE. Evolution of the cancer stem cell model. *Cell Stem Cell*. 2014;14(3):275-91.
- Krogh A. The number and distribution of capillaries in muscles with calculations of the oxygen pressure head necessary for supplying the tissue. *J Physiol*. 1919;52(6):409-15.
- Kronke J, Bullinger L, Teleanu V, Tschurtz F, Gaidzik VI, Kuhn MW, et al. Clonal evolution in relapsed NPM1-mutated acute myeloid leukemia. *Blood*. 2013;122(1):100-8.
- Kuhl C, Atzberger A, Iborra F, Nieswandt B, Porcher C, Vyas P. GATA1-mediated megakaryocyte differentiation and growth control can be uncoupled and mapped to different domains in GATA1. *Mol Cell Biol*. 2005;25(19):8592-606.
- Lambert M, Jambon S, Depauw S, David-Cordonnier MH. Targeting Transcription Factors for Cancer Treatment. *Molecules*. 2018;23(6).
- Lansdorp PM, Sutherland HJ, Eaves CJ. Selective expression of CD45 isoforms on functional subpopulations of CD34+ hemopoietic cells from human bone marrow. *J Exp Med*. 1990;172(1):363-6.
- Lapidot T, Sirard C, Vormoor J, Murdoch B, Hoang T, Caceres-Cortes J, et al. A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature*. 1994;367(6464):645-8.
- Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature*. 2013;499(7457):214-8.
- Lebert-Ghali CE, Fournier M, Kettle L, Thompson A, Sauvageau G, Bijl JJ. Hoxa cluster genes determine the proliferative activity of adult mouse hematopoietic stem and progenitor cells. *Blood*. 2016;127(1):87-90.
- Lee HK, Kim HW, Lee IY, Lee J, Lee J, Jung DS, et al. G-749, a novel FLT3 kinase inhibitor, can overcome drug resistance for the treatment of acute myeloid leukemia. *Blood*. 2014;123(14):2209-19.
- Lehnertz B, Pabst C, Su L, Miller M, Liu F, Yi L, et al. The methyltransferase G9a regulates HoxA9-dependent transcription in AML. *Genes Dev*. 2014;28(4):317-27.
- Lennartsson J, Jelacic T, Linnekin D, Shivakrupa R. Normal and oncogenic forms of the receptor tyrosine kinase kit. *Stem Cells*. 2005;23(1):16-43.
- Ley TJ, Ding L, Walter MJ, McLellan MD, Lamprecht T, Larson DE, et al. DNMT3A mutations in acute myeloid leukemia. *N Engl J Med*. 2010;363(25):2424-33.
- Li B, Carey M, Workman JL. The role of chromatin during transcription. *Cell*. 2007;128(4):707-19.
- Li Y, Vecchiarelli-Federico LM, Li YJ, Egan SE, Spaner D, Hough MR, et al. The miR-17-92 cluster expands multipotent hematopoietic progenitors whereas imbalanced expression of its individual oncogenic miRNAs promotes leukemia in mice. *Blood*. 2012;119(19):4486-98.

- Li Z, Luo RT, Mi S, Sun M, Chen P, Bao J, et al. Consistent deregulation of gene expression between human and murine MLL rearrangement leukemias. *Cancer Res.* 2009;69(3):1109-16.
- Lindsley RC, Mar BG, Mazzola E, Grauman PV, Shareef S, Allen SL, et al. Acute myeloid leukemia ontogeny is defined by distinct somatic mutations. *Blood.* 2015;125(9):1367-76.
- Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, et al. Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature.* 2009;462(7271):315-22.
- Lowenberg B, Ossenkoppele GJ, van Putten W, Schouten HC, Graux C, Ferrant A, et al. High-dose daunorubicin in older patients with acute myeloid leukemia. *N Engl J Med.* 2009;361(13):1235-48.
- Lu C, Ward PS, Kapoor GS, Rohle D, Turcan S, Abdel-Wahab O, et al. IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature.* 2012;483(7390):474-8.
- Mann G, Attarbaschi A, Schrappe M, De Lorenzo P, Peters C, Hann I, et al. Improved outcome with hematopoietic stem cell transplantation in a poor prognostic subgroup of infants with mixed-lineage-leukemia (MLL)-rearranged acute lymphoblastic leukemia: results from the Interfant-99 Study. *Blood.* 2010;116(15):2644-50.
- Marcucci G, Haferlach T, Dohner H. Molecular genetics of adult acute myeloid leukemia: prognostic and therapeutic implications. *J Clin Oncol.* 2011;29(5):475-86.
- Marcucci G, Maharry K, Wu YZ, Radmacher MD, Mrozek K, Margeson D, et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol.* 2010;28(14):2348-55.
- Marcucci G, Metzeler KH, Schwind S, Becker H, Maharry K, Mrozek K, et al. Age-related prognostic impact of different types of DNMT3A mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J Clin Oncol.* 2012;30(7):742-50.
- Maroc N, Rottapel R, Rosnet O, Marchetto S, Lavezzi C, Mannoni P, et al. Biochemical characterization and analysis of the transforming potential of the FLT3/FLK2 receptor tyrosine kinase. *Oncogene.* 1993;8(4):909-18.
- Masetti R, Vendemini F, Zama D, Biagi C, Pession A, Locatelli F. Acute myeloid leukemia in infants: biology and treatment. *Front Pediatr.* 2015;3:37.
- Mazo IB, Gutierrez-Ramos JC, Frenette PS, Hynes RO, Wagner DD, von Andrian UH. Hematopoietic progenitor cell rolling in bone marrow microvessels: parallel contributions by endothelial selectins and vascular cell adhesion molecule 1. *J Exp Med.* 1998;188(3):465-74.
- McKerrell T, Moreno T, Ponstingl H, Bolli N, Dias JM, Tischler G, et al. Development and validation of a comprehensive genomic diagnostic tool for myeloid malignancies. *Blood.* 2016;128(1):e1-9.
- Medinger M, Lengerke C, Passweg J. Novel therapeutic options in Acute Myeloid Leukemia. *Leuk Res Rep.* 2016;6:39-49.
- Medinger M, Passweg JR. Acute myeloid leukaemia genomics. *Br J Haematol.* 2017;179(4):530-42.

- Mendez-Ferrer S, Battista M, Frenette PS. Cooperation of beta(2)- and beta(3)-adrenergic receptors in hematopoietic progenitor cell mobilization. *Ann N Y Acad Sci.* 2010;1192:139-44.
- Mendler JH, Maharry K, Radmacher MD, Mrozek K, Becker H, Metzeler KH, et al. RUNX1 mutations are associated with poor outcome in younger and older patients with cytogenetically normal acute myeloid leukemia and with distinct gene and MicroRNA expression signatures. *J Clin Oncol.* 2012;30(25):3109-18.
- Meshinchi S, Alonzo TA, Stirewalt DL, Zwaan M, Zimmerman M, Reinhardt D, et al. Clinical implications of FLT3 mutations in pediatric AML. *Blood.* 2006;108(12):3654-61.
- Metzeler KH, Becker H, Maharry K, Radmacher MD, Kohlschmidt J, Mrozek K, et al. ASXL1 mutations identify a high-risk subgroup of older patients with primary cytogenetically normal AML within the ELN Favorable genetic category. *Blood.* 2011;118(26):6920-9.
- Metzeler KH, Maharry K, Radmacher MD, Mrozek K, Margeson D, Becker H, et al. TET2 mutations improve the new European LeukemiaNet risk classification of acute myeloid leukemia: a Cancer and Leukemia Group B study. *J Clin Oncol.* 2011;29(10):1373-81.
- Meyer C, Hofmann J, Burmeister T, Groger D, Park TS, Emerenciano M, et al. The MLL recombinome of acute leukemias in 2013. *Leukemia.* 2013;27(11):2165-76.
- Meyer SC, Levine RL. Translational implications of somatic genomics in acute myeloid leukaemia. *Lancet Oncol.* 2014;15(9):e382-94.
- Meyers S, Downing JR, Hiebert SW. Identification of AML-1 and the (8;21) translocation protein (AML-1/ETO) as sequence-specific DNA-binding proteins: the runt homology domain is required for DNA binding and protein-protein interactions. *Mol Cell Biol.* 1993;13(10):6336-45.
- Morita Y, Ema H, Nakauchi H. Heterogeneity and hierarchy within the most primitive hematopoietic stem cell compartment. *J Exp Med.* 2010;207(6):1173-82.
- Mrozek K, Marcucci G, Paschka P, Whitman SP, Bloomfield CD. Clinical relevance of mutations and gene-expression changes in adult acute myeloid leukemia with normal cytogenetics: are we ready for a prognostically prioritized molecular classification? *Blood.* 2007;109(2):431-48.
- Muntean AG, Hess JL. The pathogenesis of mixed-lineage leukemia. *Annu Rev Pathol.* 2012;7:283-301.
- Nio K, Yamashita T, Okada H, Kondo M, Hayashi T, Hara Y, et al. Defeating EpCAM(+) liver cancer stem cells by targeting chromatin remodeling enzyme CHD4 in human hepatocellular carcinoma. *J Hepatol.* 2015;63(5):1164-72.
- North T, Gu TL, Stacy T, Wang Q, Howard L, Binder M, et al. Cbfa2 is required for the formation of intra-aortic hematopoietic clusters. *Development.* 1999;126(11):2563-75.
- North TE, Stacy T, Matheny CJ, Speck NA, de Bruijn MF. Runx1 is expressed in adult mouse hematopoietic stem cells and differentiating myeloid and lymphoid cells, but not in maturing erythroid cells. *Stem Cells.* 2004;22(2):158-68.



- Notta F, Doulatov S, Laurenti E, Poepl A, Jurisica I, Dick JE. Isolation of single human hematopoietic stem cells capable of long-term multilineage engraftment. *Science*. 2011;333(6039):218-21.
- Nowell PC. The clonal evolution of tumor cell populations. *Science*. 1976;194(4260):23-8.
- O'Donnell MR, Abboud CN, Altman J, Appelbaum FR, Arber DA, Attar E, et al. NCCN Clinical Practice Guidelines Acute myeloid leukemia. *J Natl Compr Canc Netw*. 2012;10(8):984-1021.
- O'Shaughnessy A, Hendrich B. CHD4 in the DNA-damage response and cell cycle progression: not so NuRDy now. *Biochem Soc Trans*. 2013;41(3):777-82.
- Omatsu Y, Sugiyama T, Kohara H, Kondoh G, Fujii N, Kohno K, et al. The essential functions of adipo-osteogenic progenitors as the hematopoietic stem and progenitor cell niche. *Immunity*. 2010;33(3):387-99.
- Orkin SH. Diversification of haematopoietic stem cells to specific lineages. *Nat Rev Genet*. 2000;1(1):57-64.
- Orkin SH, Zon LI. Hematopoiesis: an evolving paradigm for stem cell biology. *Cell*. 2008;132(4):631-44.
- Osawa M, Hanada K, Hamada H, Nakauchi H. Long-term lymphohematopoietic reconstitution by a single CD34-low/negative hematopoietic stem cell. *Science*. 1996;273(5272):242-5.
- Pabst T, Eyholzer M, Haefliger S, Schardt J, Mueller BU. Somatic CEBPA mutations are a frequent second event in families with germline CEBPA mutations and familial acute myeloid leukemia. *J Clin Oncol*. 2008;26(31):5088-93.
- Palis J. Interaction of the Macrophage and Primitive Erythroid Lineages in the Mammalian Embryo. *Front Immunol*. 2016;7:669.
- Palis J, Robertson S, Kennedy M, Wall C, Keller G. Development of erythroid and myeloid progenitors in the yolk sac and embryo proper of the mouse. *Development*. 1999;126(22):5073-84.
- Papaemmanuil E, Gerstung M, Bullinger L, Gaidzik VI, Paschka P, Roberts ND, et al. Genomic Classification and Prognosis in Acute Myeloid Leukemia. *N Engl J Med*. 2016;374(23):2209-21.
- Park IK, Qian D, Kiel M, Becker MW, Pihlaja M, Weissman IL, et al. Bmi-1 is required for maintenance of adult self-renewing haematopoietic stem cells. *Nature*. 2003;423(6937):302-5.
- Parkin J, Cohen B. An overview of the immune system. *Lancet*. 2001;357(9270):1777-89.
- Paschka P, Marcucci G, Ruppert AS, Mrozek K, Chen H, Kittles RA, et al. Adverse prognostic significance of KIT mutations in adult acute myeloid leukemia with inv(16) and t(8;21): a Cancer and Leukemia Group B Study. *J Clin Oncol*. 2006;24(24):3904-11.
- Paschka P, Schlenk RF, Gaidzik VI, Habdank M, Kronke J, Bullinger L, et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J Clin Oncol*. 2010;28(22):3636-43.

- Patel JP, Gonen M, Figueroa ME, Fernandez H, Sun Z, Racevskis J, et al. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N Engl J Med*. 2012;366(12):1079-89.
- Petersdorf SH, Kopecky KJ, Slovak M, Willman C, Nevill T, Brandwein J, et al. A phase 3 study of gemtuzumab ozogamicin during induction and postconsolidation therapy in younger patients with acute myeloid leukemia. *Blood*. 2013;121(24):4854-60.
- Pina C, Enver T. Differential contributions of haematopoietic stem cells to foetal and adult haematopoiesis: insights from functional analysis of transcriptional regulators. *Oncogene*. 2007;26(47):6750-65.
- Polo SE, Kaidi A, Baskcomb L, Galanty Y, Jackson SP. Regulation of DNA-damage responses and cell-cycle progression by the chromatin remodelling factor CHD4. *EMBO J*. 2010;29(18):3130-9.
- Popel AS. Theory of oxygen transport to tissue. *Crit Rev Biomed Eng*. 1989;17(3):257-321.
- Port M, Bottcher M, Thol F, Ganser A, Schlenk R, Wasem J, et al. Prognostic significance of FLT3 internal tandem duplication, nucleophosmin 1, and CEBPA gene mutations for acute myeloid leukemia patients with normal karyotype and younger than 60 years: a systematic review and meta-analysis. *Ann Hematol*. 2014;93(8):1279-86.
- Prokocimer M, Molchadsky A, Rotter V. Dysfunctional diversity of p53 proteins in adult acute myeloid leukemia: projections on diagnostic workup and therapy. *Blood*. 2017;130(6):699-712.
- Qin YZ, Zhu HH, Jiang Q, Jiang H, Zhang LP, Xu LP, et al. Prevalence and prognostic significance of c-KIT mutations in core binding factor acute myeloid leukemia: a comprehensive large-scale study from a single Chinese center. *Leuk Res*. 2014;38(12):1435-40.
- Querfurth E, Schuster M, Kulesa H, Crispino JD, Doderlein G, Orkin SH, et al. Antagonism between C/EBPbeta and FOG in eosinophil lineage commitment of multipotent hematopoietic progenitors. *Genes Dev*. 2000;14(19):2515-25.
- Raaijmakers MH, Mukherjee S, Guo S, Zhang S, Kobayashi T, Schoonmaker JA, et al. Bone progenitor dysfunction induces myelodysplasia and secondary leukaemia. *Nature*. 2010;464(7290):852-7.
- Radulovic V, de Haan G, Klauke K. Polycomb-group proteins in hematopoietic stem cell regulation and hematopoietic neoplasms. *Leukemia*. 2013;27(3):523-33.
- Rafii S, Mohle R, Shapiro F, Frey BM, Moore MA. Regulation of hematopoiesis by microvascular endothelium. *Leuk Lymphoma*. 1997;27(5-6):375-86.
- Ranganathan P, Yu X, Na C, Santhanam R, Shacham S, Kauffman M, et al. Preclinical activity of a novel CRM1 inhibitor in acute myeloid leukemia. *Blood*. 2012;120(9):1765-73.
- Rhodes J, Hagen A, Hsu K, Deng M, Liu TX, Look AT, et al. Interplay of pu.1 and gata1 determines myelo-erythroid progenitor cell fate in zebrafish. *Dev Cell*. 2005;8(1):97-108.
- Riddell J, Gazit R, Garrison BS, Guo G, Saadatpour A, Mandal PK, et al. Reprogramming committed murine blood cells to induced hematopoietic stem cells with defined factors. *Cell*. 2014;157(3):549-64.

- Riggs AD. X inactivation, differentiation, and DNA methylation. *Cytogenet Cell Genet.* 1975;14(1):9-25.
- Rubnitz JE, Inaba H. Childhood acute myeloid leukaemia. *Br J Haematol.* 2012;159(3):259-76.
- Rubnitz JE, Inaba H, Leung WH, Pounds S, Cao X, Campana D, et al. Definition of cure in childhood acute myeloid leukemia. *Cancer.* 2014;120(16):2490-6.
- Rucker FG, Schlenk RF, Bullinger L, Kayser S, Teleanu V, Kett H, et al. TP53 alterations in acute myeloid leukemia with complex karyotype correlate with specific copy number alterations, monosomal karyotype, and dismal outcome. *Blood.* 2012;119(9):2114-21.
- Sabattini E, Bacci F, Sagranso C, Pileri SA. WHO classification of tumours of haematopoietic and lymphoid tissues in 2008: an overview. *Pathologica.* 2010;102(3):83-7.
- Sattler M, Salgia R. Targeting c-Kit mutations: basic science to novel therapies. *Leuk Res.* 2004;28 Suppl 1:S11-20.
- Saultz JN, Garzon R. Acute Myeloid Leukemia: A Concise Review. *J Clin Med.* 2016;5(3).
- Saxonov S, Berg P, Brutlag DL. A genome-wide analysis of CpG dinucleotides in the human genome distinguishes two distinct classes of promoters. *Proc Natl Acad Sci U S A.* 2006;103(5):1412-7.
- Saygin C, Carraway HE. Emerging therapies for acute myeloid leukemia. *J Hematol Oncol.* 2017;10(1):93.
- Schepers K, Pietras EM, Reynaud D, Flach J, Binnewies M, Garg T, et al. Myeloproliferative neoplasia remodels the endosteal bone marrow niche into a self-reinforcing leukemic niche. *Cell Stem Cell.* 2013;13(3):285-99.
- Schnittger S, Schoch C, Kern W, Mecucci C, Tschulik C, Martelli MF, et al. Nucleophosmin gene mutations are predictors of favorable prognosis in acute myelogenous leukemia with a normal karyotype. *Blood.* 2005;106(12):3733-9.
- Schofield R. The relationship between the spleen colony-forming cell and the haemopoietic stem cell. *Blood Cells.* 1978;4(1-2):7-25.
- Shah N, Sukumar S. The Hox genes and their roles in oncogenesis. *Nat Rev Cancer.* 2010;10(5):361-71.
- Shepherd BE, Gutter P, Lansdorp PM, Abkowitz JL. Estimating human hematopoietic stem cell kinetics using granulocyte telomere lengths. *Exp Hematol.* 2004;32(11):1040-50.
- Shikama N, Chan HM, Krstic-Demonacos M, Smith L, Lee CW, Cairns W, et al. Functional interaction between nucleosome assembly proteins and p300/CREB-binding protein family coactivators. *Mol Cell Biol.* 2000;20(23):8933-43.
- Shivarov V, Gueorguieva R, Stoimenov A, Tiu R. DNMT3A mutation is a poor prognosis biomarker in AML: results of a meta-analysis of 4500 AML patients. *Leuk Res.* 2013;37(11):1445-50.
- Shivdasani RA, Fujiwara Y, McDevitt MA, Orkin SH. A lineage-selective knockout establishes the critical role of transcription factor GATA-1 in megakaryocyte growth and platelet development. *EMBO J.* 1997;16(13):3965-73.

- Shlush LI, Zandi S, Mitchell A, Chen WC, Brandwein JM, Gupta V, et al. Identification of pre-leukaemic haematopoietic stem cells in acute leukaemia. *Nature*. 2014;506(7488):328-33.
- Small D. FLT3 mutations: biology and treatment. *Hematology Am Soc Hematol Educ Program*. 2006:178-84.
- Somervaille TC, Cleary ML. Identification and characterization of leukemia stem cells in murine MLL-AF9 acute myeloid leukemia. *Cancer Cell*. 2006;10(4):257-68.
- Spooncer E, Heyworth CM, Dunn A, Dexter TM. Self-renewal and differentiation of interleukin-3-dependent multipotent stem cells are modulated by stromal cells and serum factors. *Differentiation*. 1986;31(2):111-8.
- Starck J, Cohet N, Gonnet C, Sarrazin S, Doubeikovskaia Z, Doubeikovski A, et al. Functional cross-antagonism between transcription factors FLI-1 and EKLF. *Mol Cell Biol*. 2003;23(4):1390-402.
- Stewart MH, Albert M, Sroczynska P, Cruickshank VA, Guo Y, Rossi DJ, et al. The histone demethylase Jarid1b is required for hematopoietic stem cell self-renewal in mice. *Blood*. 2015;125(13):2075-8.
- Sudhindra A, Smith CC. FLT3 inhibitors in AML: are we there yet? *Curr Hematol Malig Rep*. 2014;9(2):174-85.
- Sugiyama T, Kohara H, Noda M, Nagasawa T. Maintenance of the hematopoietic stem cell pool by CXCL12-CXCR4 chemokine signaling in bone marrow stromal cell niches. *Immunity*. 2006;25(6):977-88.
- Sun XJ, Man N, Tan Y, Nimer SD, Wang L. The Role of Histone Acetyltransferases in Normal and Malignant Hematopoiesis. *Front Oncol*. 2015;5:108.
- Super HJ, McCabe NR, Thirman MJ, Larson RA, Le Beau MM, Pedersen-Bjergaard J, et al. Rearrangements of the MLL gene in therapy-related acute myeloid leukemia in patients previously treated with agents targeting DNA-topoisomerase II. *Blood*. 1993;82(12):3705-11.
- Tachibana M, Ueda J, Fukuda M, Takeda N, Ohta T, Iwanari H, et al. Histone methyltransferases G9a and GLP form heteromeric complexes and are both crucial for methylation of euchromatin at H3-K9. *Genes Dev*. 2005;19(7):815-26.
- Takei H, Kobayashi SS. Targeting transcription factors in acute myeloid leukemia. *Int J Hematol*. 2018.
- Tang JL, Hou HA, Chen CY, Liu CY, Chou WC, Tseng MH, et al. AML1/RUNX1 mutations in 470 adult patients with de novo acute myeloid leukemia: prognostic implication and interaction with other gene alterations. *Blood*. 2009;114(26):5352-61.
- Taoudi S, Bee T, Hilton A, Knezevic K, Scott J, Willson TA, et al. ERG dependence distinguishes developmental control of hematopoietic stem cell maintenance from hematopoietic specification. *Genes Dev*. 2011;25(3):251-62.
- Thomas D, Majeti R. Biology and relevance of human acute myeloid leukemia stem cells. *Blood*. 2017;129(12):1577-85.
- Tiley S, Claxton D. Clofarabine in the treatment of acute myeloid leukemia in older adults. *Ther Adv Hematol*. 2013;4(1):5-13.

- Tober J, Koniski A, McGrath KE, Vemishetti R, Emerson R, de Mesy-Bentley KK, et al. The megakaryocyte lineage originates from hemangioblast precursors and is an integral component both of primitive and of definitive hematopoiesis. *Blood*. 2007;109(4):1433-41.
- Trowbridge JJ, Orkin SH. Dnmt3a silences hematopoietic stem cell self-renewal. *Nat Genet*. 2011;44(1):13-4.
- Trowbridge JJ, Snow JW, Kim J, Orkin SH. DNA methyltransferase 1 is essential for and uniquely regulates hematopoietic stem and progenitor cells. *Cell Stem Cell*. 2009;5(4):442-9.
- Vardiman JW, Thiele J, Arber DA, Brunning RD, Borowitz MJ, Porwit A, et al. The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: rationale and important changes. *Blood*. 2009;114(5):937-51.
- Velten L, Haas SF, Raffel S, Blaszkiewicz S, Islam S, Hennig BP, et al. Human haematopoietic stem cell lineage commitment is a continuous process. *Nat Cell Biol*. 2017;19(4):271-81.
- Velu CS, Baktula AM, Grimes HL. Gfi1 regulates miR-21 and miR-196b to control myelopoiesis. *Blood*. 2009;113(19):4720-8.
- Visnjic D, Kalajzic Z, Rowe DW, Katavic V, Lorenzo J, Aguila HL. Hematopoiesis is severely altered in mice with an induced osteoblast deficiency. *Blood*. 2004;103(9):3258-64.
- Voigt P, Tee WW, Reinberg D. A double take on bivalent promoters. *Genes Dev*. 2013;27(12):1318-38.
- Waddington CH. The epigenotype. 1942. *Int J Epidemiol*. 2012;41(1):10-3.
- Walmsley M, Ciau-Uitz A, Patient R. Adult and embryonic blood and endothelium derive from distinct precursor populations which are differentially programmed by BMP in *Xenopus*. *Development*. 2002;129(24):5683-95.
- Walter RB, Othus M, Paietta EM, Racevskis J, Fernandez HF, Lee JW, et al. Effect of genetic profiling on prediction of therapeutic resistance and survival in adult acute myeloid leukemia. *Leukemia*. 2015;29(10):2104-7.
- Wang CQ, Krishnan V, Tay LS, Chin DW, Koh CP, Chooi JY, et al. Disruption of Runx1 and Runx3 leads to bone marrow failure and leukemia predisposition due to transcriptional and DNA repair defects. *Cell Rep*. 2014;8(3):767-82.
- Wang H, La Russa M, Qi LS. CRISPR/Cas9 in Genome Editing and Beyond. *Annu Rev Biochem*. 2016;85:227-64.
- Wang JC, Doedens M, Dick JE. Primitive human hematopoietic cells are enriched in cord blood compared with adult bone marrow or mobilized peripheral blood as measured by the quantitative in vivo SCID-repopulating cell assay. *Blood*. 1997;89(11):3919-24.
- Ward PS, Patel J, Wise DR, Abdel-Wahab O, Bennett BD, Collier HA, et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell*. 2010;17(3):225-34.
- Wikipedia C. Hematopoietic stem cell. 2018.

- Wilkerson MJ. Principles and applications of flow cytometry and cell sorting in companion animal medicine. *Vet Clin North Am Small Anim Pract.* 2012;42(1):53-71.
- Wilkes MC, Repellin CE, Sakamoto KM. Beyond mRNA: The role of non-coding RNAs in normal and aberrant hematopoiesis. *Mol Genet Metab.* 2017;122(3):28-38.
- Wilson A, Laurenti E, Oser G, van der Wath RC, Blanco-Bose W, Jaworski M, et al. Hematopoietic stem cells reversibly switch from dormancy to self-renewal during homeostasis and repair. *Cell.* 2008;135(6):1118-29.
- Winkler IG, Sims NA, Pettit AR, Barbier V, Nowlan B, Helwani F, et al. Bone marrow macrophages maintain hematopoietic stem cell (HSC) niches and their depletion mobilizes HSCs. *Blood.* 2010;116(23):4815-28.
- Winters AC, Bernt KM. MLL-Rearranged Leukemias-An Update on Science and Clinical Approaches. *Front Pediatr.* 2017;5:4.
- Witko-Sarsat V, Rieu P, Descamps-Latscha B, Lesavre P, Halbwachs-Mecarelli L. Neutrophils: molecules, functions and pathophysiological aspects. *Lab Invest.* 2000;80(5):617-53.
- Wouters BJ, Lowenberg B, Erpelinck-Verschueren CA, van Putten WL, Valk PJ, Delwel R. Double CEBPA mutations, but not single CEBPA mutations, define a subgroup of acute myeloid leukemia with a distinctive gene expression profile that is uniquely associated with a favorable outcome. *Blood.* 2009;113(13):3088-91.
- Xiao P, Dolinska M, Sandhow L, Kondo M, Johansson AS, Boudierlique T, et al. Sip1 deficiency-induced bone marrow niche alterations lead to the initiation of myeloproliferative neoplasm. *Blood Adv.* 2018;2(5):534-48.
- Yamamoto JF, Goodman MT. Patterns of leukemia incidence in the United States by subtype and demographic characteristics, 1997-2002. *Cancer Causes Control.* 2008;19(4):379-90.
- Yamamoto R, Morita Y, Ooehara J, Hamanaka S, Onodera M, Rudolph KL, et al. Clonal analysis unveils self-renewing lineage-restricted progenitors generated directly from hematopoietic stem cells. *Cell.* 2013;154(5):1112-26.
- Yoshihara H, Arai F, Hosokawa K, Hagiwara T, Takubo K, Nakamura Y, et al. Thrombopoietin/MPL signaling regulates hematopoietic stem cell quiescence and interaction with the osteoblastic niche. *Cell Stem Cell.* 2007;1(6):685-97.
- Yu MG, Zheng HY. Acute Myeloid Leukemia: Advancements in Diagnosis and Treatment. *Chin Med J (Engl).* 2017;130(2):211-8.
- Yu VW, Saez B, Cook C, Lotinun S, Pardo-Saganta A, Wang YH, et al. Specific bone cells produce DLL4 to generate thymus-seeding progenitors from bone marrow. *J Exp Med.* 2015;212(5):759-74.
- Yu VW, Scadden DT. Hematopoietic Stem Cell and Its Bone Marrow Niche. *Curr Top Dev Biol.* 2016;118:21-44.
- Zhang J, Niu C, Ye L, Huang H, He X, Tong WG, et al. Identification of the haematopoietic stem cell niche and control of the niche size. *Nature.* 2003;425(6960):836-41.
- Zhang P, Iwasaki-Arai J, Iwasaki H, Fenyus ML, Dayaram T, Owens BM, et al. Enhancement of hematopoietic stem cell repopulating capacity and self-renewal in the absence of the transcription factor C/EBP alpha. *Immunity.* 2004;21(6):853-63.

- Zhang X, Lian Z, Padden C, Gerstein MB, Rozowsky J, Snyder M, et al. A myelopoiesis-associated regulatory intergenic noncoding RNA transcript within the human HOXA cluster. *Blood*. 2009;113(11):2526-34.
- Zhang X, Lv D, Zhang Y, Liu Q, Li Z. Clonal evolution of acute myeloid leukemia highlighted by latest genome sequencing studies. *Oncotarget*. 2016;7(36):58586-94.
- Zhou J, Bi C, Janakakumara JV, Liu SC, Chng WJ, Tay KG, et al. Enhanced activation of STAT pathways and overexpression of survivin confer resistance to FLT3 inhibitors and could be therapeutic targets in AML. *Blood*. 2009;113(17):4052-62.
- Zhu J, Garrett R, Jung Y, Zhang Y, Kim N, Wang J, et al. Osteoblasts support B-lymphocyte commitment and differentiation from hematopoietic stem cells. *Blood*. 2007;109(9):3706-12.
- Zovein AC, Hofmann JJ, Lynch M, French WJ, Turlo KA, Yang Y, et al. Fate tracing reveals the endothelial origin of hematopoietic stem cells. *Cell Stem Cell*. 2008;3(6):625-36.